THE ZOOLOGICAL RECORD

Museuma

Z 7991 , 287 v. 96

no. 5

VOLUME 96 SECTION 5 1959

ECHINODERMATA

COMPILED BY

A. M. CLARK, M.A.

LONDON
PUBLISHED BY
THE ZOOLOGICAL SOCIETY OF LONDON

PRICE FOUR SHILLINGS July 1961



It will greatly help in the production of The Zoological Record and assist the Recorders of the individual sections, if authors will forward a copy of their paper or memoir to the Editor of the Zoological Record, The Zoological Society of London, Regent's Park, London, N.W.1. In the case of separately printed copies of articles so forwarded, the original pagination should be given.

All business correspondence concerning the Record should be addressed to the Scientific Director, The Zoological Society of London, Regent's Park, London, N.W.1.

THE MUSEUMS JOURNAL

Published by the Museums Association 22 Fitzroy Street, Fitzroy Square, London, W.1.

This journal, which is published monthly, contains articles, reviews and technical notes on every Museum aspect of zoology and other sciences. It is the only English journal which deals with the installation and preservation of exhibits, and which reviews Museum work in all parts of the world.

It is published monthly price 4s., by the Association and can be obtained from the above address. It is distributed free to all members of the Museums Association (Subscription £3 3s. 0d. per annum), of which full particulars are obtainable from the Secretary at the above address.

ECHINODERMATA

COMPILED BY

A. M. CLARK, M.A.

FI THE PURPLE STATE

ECHINODERMATA

CONTENTS

I.	TITLES	•••	***	***	***	***	 ***		***	1
II.	SUBJECT IN	DEX				***	 	***		9
III.	SYSTEMATIC	INDEX					 			12

I.—TITLES

In the list of titles the year has been omitted where it is the same as the volume year of the "Record", namely (1959).

Atzelius, B. Electron microscopy of the sperm tail. Results obtained with a new fixative. J. Biophys. & Biochem. Cytol. 5: 269-278 2 pls. 1 fig.

Agrell, I. (1). The structure of the chromatine in the pronucleus and the syncarion of the sea urchin egg. Ark. Zool. 12:95-98 l pl. l fig.

Agrell, I. (2). The influence of the temperature upon the nuclear changes during syngamy in the sea urchin egg. Ark. Zool. 12: 291-300 2 pls.

Aketa, K. & Tomita, H. Oxidative phosphorylation by a cell-free particulate system from unfertilized eggs of the sea urchin, *Hemicentrotus pulcherrimus*. Bull. mar. biol. Sta. Asamushi 9 1958: 57-58.

Alcals, A. C. Sea urchin eggs: their use in the laboratory with notes on *Tripneustes gratilla* (Linn.). Silliman J. 4 1957: 192–195 3 pls.

Allen, R. D. see Roslansky, J. D.

Allen, R. D. & Bunim, L. Changes in cytoplasmic structure following fertilization in *Arbacia* eggs. Biol. Bull. Woods Hole 117: 403.

Ancona-Lopez, A. A. Sôbre holotúrias do litoral sul Brasileiro. Bol. Fac. Filos. Cienc. Univ. S. Paulo, no. 223 Zool. 21 1958: 1-53. [Portuguese with English summary.]

Anderson, J. M. (1). Regeneration of pyloric caeca in starfishes. Anat. Rec. 134: 526.

Anderson, J. M. (2). Studies on the cardiac stomach of a starfish, *Patiria miniata* (Brandt). Biol. Bull. Woods Hole 117: 185-201 21 figs.

Ando, S. On fucose in the sea urchin egg. Bull. mar. biol. Sta. Asamushi 9 1958: 55-56 2 figs.

Andrade, M. M. de & Andrade, J. B. M. de. Estado actual dos conhecimentos sobre a paleontologia de Angola (até fins de 1955). An. Jta. Inv. Ultramar, Lisboa 12 Tom. 4 1957: 5-209.

Bäckström, S. (1). Changes in ribonucleic acid content during early sea urchin development. Ark. Zool. 12: 339–342.

Bäckström, S. (2). Studies on sulphydryl-containing substances in sea urchin embryos of various developmental trends. Exper. Cell Res. 16: 165-173 3 figs.

Badillo, L. Catalogo de especies fosiles del Museo del Instituto Geologico y Minero de España. Notas Inst. geol. Esp. 55: 71-124 50 pls. (Crinoid on p. 88).

Balavoins, P. Sur la présence de Semiometra italica (Schlüter, 1878) (Comatulidae) dans le Lutétien de Bois-Gonët (Loire-Atlantique). Bull. Mus. Hist. nat. Paris (2) 30: 525.

Balinsky, B. I. An electron microscopic investigation of the mechanisms of adhesion of the cells in a sea urchin blastula and gastrula. Exper. Cell Res. 16: 429-433 9 figs.

Baltus, E. see Vincent, W. S.

Bang, F. B. & Chast, A. B. The effect of starfish toxin on amebocytes. Biol. Bull. Woods Hole 117: 403-404.

Bernstein, G. S. & Black, R. E. Inhibition of cleavage and development of the sea-urchin egg by 2-desoxy-D-glucose. Proc. Soc. exp. Biol. Med. 102:531-534.

Bernstein, M. H. Functional modifications of sperm structure. Biol. Bull. Woods Hole 117:433.

Bismarck, O. von. Versuch einer Analyse der die Stoffwechselintensität ("Ruheumsatz") von Asterias rubens L. beeinflussenden Faktoren. Kieler Meeresforsch. 15: 164–186 9 pls.

Black, R. E. & Tyler, A. (1). Effects of fertilization and development on the oxidation of carbon monoxide by eggs of Strongylocentrotus and Urechis as determined by use of C¹³. Biol. Bull. Woods Hole 117: 443-453.

Black, R. E. see Bernstein, G. S.

Black, R. E. & Tyler, A. (2). Cytochrome oxidase and oxidation of CO in eggs of the sea urchin Strongylocentrotus purpuratus. Biol. Bull. Woods Hole 117: 454-457.

Boolootian, R. A. A simple technique for studying in vivo skeletal structures of echinoderm larvae. Turtox News 37: 115.

Boolootian, R. A. & Giese, A. C. Clotting of echinoderm coelomic fluid. J. exp. Zool. 140: 207-229 3 pls.

Boolootian, R. A. & others. A contribution to the biology of a deep sea echinoid, Allocentrotus fragilis (Jackson). Biol. Bull. Woods Hole 116: 362-372 3 figs.

Booloctian, R. A. & Moore, A. R. A case of ovotestes in the sea urchin Strongylocentrotus purpuratus. Science 129: 271-272.

Boström, H. see Perlmann, P.

Bougis, P. Sur l'effet biologique del cuivre en eau de mer. CR Acad. Sci. Paris 249: 326-328 l fig. Branham, J. M. & Metz, C. B. Inhibition of fertilization and agglutination in *Arbacia* by an extract of *Fucus*. Biol. Bull. Woods Hole 117: 392–393.

Brice, M. C. (1). Toxicité des colorants metachromatiques pour l'oeuf d'Etoile de mer. Arch. Biol. Paris 70: 121-125.

Brice, M. C. (2). Animalisation et végétalisation dans des oeufs d'oursin soumis à des colorants métachromatiques. Arch. Biol. Paris 70: 127-132 pl. v.

Brito, I. M. Sôbre uma nova *Clypeaster* do Brasil (Echinoidea Clypeasteroidea). Avulso Centr. Est. zool. Univ. Brasil No. 1: 1-4 2 pls.

Brookbank, J. W. (1). The respiration of unfertilized sea urchin eggs in the presence of antisera against fertilizin. Biol. Bull. Woods Hole 116: 217-225.

Brookbank, J. W. (2). The presence of fertilizin haptens within the unfertilized sea urchin egg. Biol. Bull. Woods Hole 117: 63-67.

Brooks, M. M. Negative oxidation reduction potential as the limiting factor for exogastrulation in sea urchins. Protoplasma **51**: 131–153.

Bullivant, J. S. Photographs of the bottom fauna in the Ross Sea. N.Z. J. Sci. 2: 485-497 10 figs.

Bunim, L. see Allen, R.D.

Byers, T. F. see Heilbrunn, L. V.

Cabibel, J., Termier H. & Termier, G. Les échinodermes mésocambriens de la montagne Noire (Sud de la France). Ann. Paléont. 44: 279-294 2 pls. 10 figs.

Chaet, A. B. & McConnaughy, R. A. Physiologic activity of nerve extracts. Biol. Bull. Woods Hole 117: 407-408.

Chaet, A. B. & Mosher, C. G. Apparent loss of biological activity of the toxin(s) from scalded starfish. Anat. Rec. 132: 421.

Chaet, A. B., see Bang, F. B.

Cheney, R. H. & Speidel, C. C. Relation of x-ray dosage to modifications in fertilization and early development of *Arbacia punctulata*. Biol. Bull. Woods Hole 117: 408.

Cheney, R. H. see Speidel, C. C.

Cherbonnier, G. (1). Echinodermes de la Guyane française (Crinoides, Astérides, Ophiurides, Echinides, Holothurides). (Notes 1-5). Bull. Mus. Hist. nat. Paris 31: 105-111, figs. 1, 2; 168-172; figs. 3, 4; 261-265, figs. 5-7; 367-372 figs. 8, 9; 440-447 figs. 10, 11.

Cherbonnier, G. (2). XXXII. Holothurioidea. Rés. Sci. Miss. R. P. Dollfus en Egypte 1927-29 Paris No. 32: 249-251.

Cherbonnier, G. (3). Échinides. Rés. Sci. Exped. Océanogr. Belge Eaux Cotières Afr. Atlant. Sud. 3 6: 35-59 10 pls. 1 fig.

Clark, D. L. Texas Cretaceous ophiuroids. J. Paleont. 33: 1126-1127.

Clark, H. E. see Fell, H. B.

Cognetti, G. see Delavault, R.

Cooke, C. W. Cenozoic echinoids of eastern United States. Prof. Pap. U.S. geol. Surv. 321: 1-106 43 pls.

Costello, D. P. A general theory of membrane elevation in marine eggs. Biol. Bull. Woods Hole 117: 410-411.

Cousineau, G. see Gross, P. R.

Cramer, H. R. Devonian starfish from Pike County, Pennsylvania. J. Paleont. 33: 471-473 1 fig.

Crisp, D. J. & Fischer-Piette, E. Répartition des principales espèces intercotidales de la côte Atlantique française en 1954–1955. An. Inst. océanogr. Paris N.S. 36: 275–387 21 figs.

Czihak, G. Vorkommen und Lebensweise de Ophiothrix quinquemaculata in de nördlichen Adria bei Rovinj. Thalassia jugosl. 16-10: 19-27 3 figs.

Dan, K. see Sakai, H.

Defiandre-Rigaud, M. Sur quelques sclérites d'Holothurides de l'Oligocène d'Innien, Holstein. Rev. Micropaléont. 1: 190-200 3 pls.

Deichmann, E. (1). Ekman's Barrier and the Holothurians of the Panama region. Proc. int. Congr. Zool. 15: 270-272.

Deichmann, E. (2). The Holothurian fauna of Antarctica. Proc. int. Congr. Zool. 15: 335-337.

Delavault, R. Premières observations sur le cycle génital d' Asterina gibbosa Pennant de Dinard. Bull. Lab. marit. Dinard No. 45: 42-44.

Delavault, R. & Cognetti, G. Recherches sur la sexualité des Asterides de la Méditerranée. Proc. int. Congr. Zool. 15: 284-286.

Domantay, J. S. (1). The zoogeographical distribution of the Indo-Pacific littoral Holothurioidea. Proc. 8th Pac. sci. Congr. 3 1957: 417-455 1 pl.

Domantay, J. S. (2). Some holothurians from Florida in the collection of the Allan Hancock Foundation. Agra Univ. J. Res. Sci. 7 1958: 181-202.

[Dubatolowa, Y. A. & Shao, T.] [Stalks of sea-lilies from Carboniferous, Permian and Triassic deposits in southern China.] Acta palaeont. Sinica 7: 41-79 2 pls. 4 figs. [In Chinese and Russian.]

Engel, H. [in] Rumphius memorial volume. Chapter 9.
The Echinoderms of Rumphius. Amsterdam pp. 209–223
4 pls.

Ennor, A. H. see Griffiths, D. E.

Eppley, R. W. & Lasker, R. Alginase in the sea urchin Strongylocentrotus purpuratus. Science 129: 214-215 2 figs.

Farmanfarmaian, A. The respiratory surface of the purple sea urchin (Strongylocentrotus purpuratus). Anat. Rec. 134: 561.

Faust, R. G. & others. Isolation and characterization of cortical granule-hyaline material of the *Arbacia* egg. Biol. Bull. Woods Hole 117: 394.

Feder, H. M. The food of the starfish, *Pisaster ochraceus*, along the California coast. Ecology **40**: 721–724 2 figs.

Fell, H. B. Starfishes of New Zealand. Tuatara 7: 127-142 3 pls.

Fell, H. B. & Clark, H. E. Anareaster, a new genus of Asteroidea from Antarctica. Trans. roy. Soc. N.Z. 87: 185–187 3 figs.

Ferreira Soares, A. (1). Sobre alguns fosseis da região de entre Lobito e Catumbela (Angola). Mem. Mus. min. geol. Univ. Coimbra 46: 11-22 5 pls. [Portuguese with French and English summaries.]

Ferreira Soares, A. (2). Contribuição para o estudo fauna fóssil da região de Entre Lobito e Catumbela (Angola). Garcia de Orta 7: 135-154 5 pls. [Portuguese with English summary.]

Fischer-Piette, E. see Crisp, D. J.

Forster, G. R. The ecology of *Echinus esculentus L.* quantitative distribution and rate of feeding. J. mar. Biol. Ass. U.K. 38: 361–367 2 pls. 2 figs.

Fukushi, T. (1). The presumptive position of the dorsal and ventral areas in the sea urchin eggs studied with local vital staining. Bull. mar. biol. Sta. Asamushi 9 1958: 75-76 1 fig.

Fukushi, T. (2). The presumptive position of the dorsal and the ventral areas in the sea urchin egg studied with local vital-staining. Bull. mar. biol. Sta. Asamushi 9: 127-132 4 figs.

Fukushi, T. (3). On the cell mass observed on the lett side of the pluteus of the sea urchin, *Temnopleurus* hardwickii. Bull. mar. biol. Sta. Asamushi 9: 133-135 4 figs.

Gautier, Y. V. Sur quelques cas d'epibiose: Bryozoaires sur *Leptometra*. Rec. Trav. Sta. mar. Endoume No. 26: 143-148.

Giese, A. C. Incidence of Syndesmis in the gut of two species of sea urchins. Anat. Rec. 132: 441-442.

Giese, A. C. see Boolootian, R. A.

Giese, A. C. & others. Organic productivity in the reproductive cycle of the purple sea urchin. Biol. Bull. Woods Hole 116: 49-58.

Gillett, K. & McNeill, F. The Great Barrier Reef and adjacent isles. Sydney 194 pp. 161 pls. (Echinodermata on pp. 95-103).

Golikova, M. N. see Poliansky, G. I.

Griffiths, D. E., Morrison, J. F. & Ennor, A. H. The distribution of guanidines, phosphagens and N-amidino phosphokinases in echinoids. Biochem. J. 65 1957: 612– 617.

Gross, P. R. & Cousineau, G. Glucose-6-phosphatase in sea urchin eggs. Biol. Bull. Woods Hole 117: 413.

Gross, P. R. & others. The block to cell division produced by deuterium oxide. Biol. Bull. Woods Hole 117: 395.

Gutschick, R. C. Lower Mississippian holothurian sclerites from the Rockford Limestone of Northern Indiana. J. Paleont. 33: 130-137 2 pls. 6 figs.

Hagström, B. E. (1). The influence of jelly coat solution on sea urchin spermatozoa. Exper. Cell Res. 16: 184-192 2 figs.

Hagström, B. E. (2). Further experiments on jelly-free sea urchin eggs. Exper. Cell Res. $17:256-261\ 1$ fig.

Hagström, B. E. (3). Experiments on hybridization of sea urchins. Ark. Zool. 12: 127-135 4 figs.

Hagström, B. E. & Hagström, B. The effect of decreased and increased temperatures on fertilization. Exper. Cell Res. 16: 174–183 3 figs.

Hagström, B. E. & Runnström, J. Refertilization of partially fertilized sea urchin eggs. Exper. Cell Res. 16: 309-314.

Halstead, B. W. Dangerous marine animals. Cambridge, Maryland. pp. [xiv] 146: frontis. col. 86 figs. (Echinoids on pp. 49–53).

Hampton, J. S. Statistical analysis of holothurian sclerites. Micropaleontology 5:335-349 4 pls. 3 figs.

Hancock, D. A. Enemies of oysters in Britain. New Scient. 5:172-174 4 figs.

Hathaway, R. R. The effects of sperm on S³⁵ labelled Arbacia fertilizin. Biol. Bull. Woods Hole 117: 395-396.

Hattin, D. E. An occurrence of Ophiuraster burrisi Miller. J. Paleont. 33: 1125-1126 1 fig.

Heilbrunn, L. V. The action of glycerol on protoplasm. Biol. Bull. Woods Hole 117: 388–389.

Heilbrunn, L. V. & Byers, T. F. A study of membrane elevation in various marine eggs. Biol. Bull. Woods Hole 117: 396–397.

Hersch, R. T. see Kane, R. E.

Hickman, J. L. Dendrogaster tasmaniensis sp. nov. (Ascothoracida) from the sea-star Allostichaster polyplax (Müller and Troschel). Parasitology 49: 316-329 19 figs.

Hiramoto, Y. (1). Preliminary report on the electric properties of echinoderm eggs. Bull. mar. biol. Sta. Asamushi 9 1958: 69-70 2 figs.

Hiramoto, Y. (2). Changes in electric properties upon fertilization in the sea urchin egg. Exper. Cell Res. 16: 421-424 2 figs.

Hori, R. On the change in membrane potential accompanying fertilization of the narcotized egg of the sea urchin, Hemicentrotus pulcherrimus. Bull. mar. biol. Sta. Asamushi 9 1958: 67-68 1 fig.

Hsiao, S. C. The source and process of uptake of calcium in the early formation of sea urchin plutei skeleton. Anat. Rec. 134: 582-583.

Hurley, D. E. Some features of the benthic environment in Cook Strait. N.Z. J. Sci. 2: 137-147 5 figs.

Immers, J. & Runnström, J. The effect of exposure of the sea urchin egg to cyanide in the early postfertilization period. Ark. Zool. 12: 83-94 10 figs.

Ishibashi, S. Studies on the muscular extracts of the lower animals. II. Muscular extract of the sea-cucumber (Stichopus japonica). Sci. Rep. Saikyo Univ., Nat. Sci. 11953: 74-76. [In Japanese.]

Ishikawa, M. Experimental formation of the mark of sperm-entry and its relation to toughening of the fertilization membrane in the sea urchin egg. Bull. mar. biol. Sta. Asamushi 9 1958: 51-52.

Jakowska, S. Coney Island sea-urchins. Anim. Kingd. 62: 137-139 2 figs.

Jakowska, S. & others. Hemopoietic effects of Holothurin, a steroid saponin from the sea-cucumber, Actinopyga agassizi, in Rana pipiens. Anat. Rec. 132: 459.

Jeannet, A. Sur quelques Echinides tertiaires de la Trinité. Mitt. naturf. Ges. Basel 70: 193-204 9 pls.

Jesionek-Szymanska, W. Remarks on the structure of the apical system of irregular echinoids. Acta palaeont. polon. 4:339-352 1 pl. 3 figs.

Kane, R. E. & Hersh, R. T. The isolation and preliminary characterization of a major soluble protein of the sea urchin egg. Exper. Cell Res. 16: 59-69 7 figs.

Kenny, R. A new Australian record of an Ascothoracid parasite. Aust. J. Sci. 21: 221.

Kinoshita, S. On the mechanism of the action of the 'relaxing factor' on the movement of the flagellum-model of the spermatozoa in some marine forms. Bull. mar. biol. Sta. Asamushi 9 1958: 59.

Kitasume, Y. Effects of Na-salicylate upon the cleavage of sea urchin eggs. Mem. Coll. Sci. Univ. Kyoto 26B 1: 15-26 4 figs.

Kitching, J. A. & others. The ecology of Lough Inc. VIII. Mussels and their predators J. Anim. Ecol. 23: 331-341 4 figs.

Kobayashi, N. see Tahara, Y.

Koenig, J. W. & Niewoehner, W. Pentececrinus, a new microcrinoid from the Louisiana formation of Missouri. J. Paleont. 33: 462-470 3 figs.

Kohler, K. & Metz, C. B. (1). Soluble and "insoluble" antigens of the *Arbacia* sperm surface. Anat. Rec. 134: 595.

Kohler, K. & Metz, C. B. (2). Extraction of egg jelly precipitating agents from *Arbacia* sperm. Biol. Bull. Woods Hole 117: 416.

Kojima, M. K. On the rhythmic formation of clear cytoplasmic spots in relation to the cleavage cycles of the sea urchin egg. Bull. mar. biol. Sta. Asamushi 9 1958: 61-82.

[Kotetishvili, E. V.] [Stratigraphy of the Cretaceous deposits of the Shkmerskoi syncline.] 40 pp. 4 pls. Tiflis 1958. [In Georgian with Russian summary.]

Kubo, K. Studies on the systematic serology of seastars. II, III and IV. Annot. zool. jap. 31 1958: 97–102, 2 figs. 32 1959: 74–80 2 figs. 214–219 1 fig.

Lallier, R. (1). Les effets de la phénazone sur la détermination embryonnaire de l'oeuf de l'oursin Paracentrotus lividus. C.R. Acad. Sci. Paris 248: 1416-1419.

Lallier, R. (2). Les groupes sulfhydriles et la détermination embryonnaire de l'oeuf de l'oursin *Paracentrotus* lividus. C.R. Acad. Sci. Paris 248: 2397–2399.

Lallier, R. (3). Recherches sur l'animalisation de l'œuf d'oursin par les ions zinc. J. Embryol. exp. Morph. 7:540-548 4 figs.

Lallier, R. (4). Végétativisation de l'œuf de l'oursin par la Phénazone. Experientia 15 : 228-229.

Lami, R. Paracentrotus lividus Lam. à Fréhel. Bull. Lab. marit. Dinard No. 45: 70.

Lang, W. D. Report on Dorset natural history for 1958: Invertebrates other than Arthropods. Proc. Dorset nat. Hist. Soc. 80: 23-24.

Lasker, R. see Eppley, R. W.

Leischner, W. Zur Mikrofazies Kalkalpiner Gesteine. S.B. öst. Akad. Wiss, Math.-nat. Kl. Abt. I. 168: 839–882 3 pls. 17 figs.

Le Maître, D. Contribution à l'étude des faunes dévoniennes d'Afrique du Nord. 1. Échinodermes. Bull. Carte géol. Alger N.S. 20 1958 : 113–154 3 pls. 3 figs.

Leone, V. The influence of ribonuclease on the development of the sea urchin. Biol. Bull. Woods Hole 117: 434-435.

Light, S. F. & others. Intertidal invertebrates of the central California coast. Berkeley $1954:pp.\ xiv, 446:138$ figs. (Echinodermata on pp. 285–294).

Litchfield, J. B. & Whiteley, A. H. Studies on the mechanism of phosphate accumulation by sea urchin embryos. Biol. Bull. Woods Hole 117: 133-149 7 figs.

Lenning, S. Variations in the thermal sensitivity during the early development of the sea urchin embryo. Ark. Zool. 12:359-381.

Lynch, W. F. Factors influencing metamorphosis of larvae of some of the sessile organisms. Proc. int. Congr. Zool. 15: 239-241.

McConnaughy, R. A. see Chaet, A. B.

McIntyre, A. D. The ecology of Scottish inshore fishing grounds. 1. The bottom fauna of east coast grounds. Mar. Res. 1958 No. 1: 1-24 3 figs.

McNeill, F. see Gillett, K.

McRae, A. Evechinus chloroticus (Val.) an endemic New Zealand echinoid. Trans. roy. Soc. N.Z. **86** 3-4: 205-267 2 pls. 15 figs.

Madsen, F. J. (1). On a new North-Atlantic sea-star, Chondraster hermanni n.sp., with some remarks on related forms. Vidensk. Medd. naturh. Foren. Kbh. 121: 153-160 4 figs.

Madsen, F. J. (2). On some Linnéan and Müllerian types of Echinoderms in the Zoological Museum of Copenhagen. Vidensk. Medd. naturh. Foren. Kbh. 121: 161–170 6 figs.

Maggio, R. Cytochrome oxidase activity in the mitochondria of unfertilized and fertilized sea urchin eggs. Exper. Cell Res. 16: 272-278 1 fig.

Manwell, C. Oxygen equilibrium of *Cucumaria* miniata hemoglobin and the absence of the Bohr effect. J. cell. comp. Physiol. 53: 75-83 1 fig.

Marsh, J. B. A rapid method for measuring total lipid, applied to the analysis of the lipids and lipoproteins of *Arbacia* eggs and embryos. Biol. Bull. Woods Hole 117: 435.

Marsland, D. Nuclear and cytoplasmic factors in cleavage induction. Anat. Rec. 132: 473.

Mattisson, A. G. M. Cytochrome c, cytochrome oxidase, and respiratory intensity in some types of invertebrate muscles. Ark. Zool. 12: 143-163 4 figs.

Menkin, V. Nature of factors regulating the rate of cell division. J. exp. Zool. 140: 441-470 7 figs.

Menkin, V. & others. Accelerator and retarding cell division factors of one species on ova of an unrelated form. J. exp. Zool. 140: 471-491.

Merriam, R. W. The origin and fate of annulate lamellae in maturing sand dollar eggs. J. Biophys. & Biochem. Cytol. 5: 117-122 5 pls.

Merson, G. The localization of some dehydrogenases in Echinarachnius parma. Biol. Bull. Woods Hole 117: 420.

Mets, C. B. (1). Inhibition of fertilizin agglutination of sperm by the dermal secretion from *Arbacia*. Biol. Bull. Woods Hole 116: 472–483 1 fig.

Metz, C. B. (2). Studies on the fertilization inhibiting action of *Arbaca* dermal secretion. Biol. Bull. Woods Hole 117: 398.

Metz, C. B. see Branham, J. M.

Metz, C. B. see Kohler, K.

Millott, N. & Yoshida, M. The photosensitivity of the sea urchin Diadema antillarum Philippi: responses to increases in light intensity. Proc. zool. Soc. Lond. 133: 67-71 1 fig. 1 pl.

Millott, N. see Yoshida, M.

[Mitrović-Petrović, J.] [Contribution à la Connaissance de la faune Crétacée Inférieure des environs de Krivi Vir (Serbie Orientale).] Ann. géol. Pén. balkan. 25 1958 : 43-60 3 pls. [In Russian with French summary on p. 60.1

Mohri, H. Change in acyl-ester content of lipid fraction in sea-urchin spermatozoa during aging. Annot. Zool. jap. 32:170-173.

Monroy, A. & Nakano, E. Evaluation of the methods for the incorporation of radioactive compounds in the echinoderm eggs. Pubbl. Staz. zool. Napoli 31 Suppl.: 95-99. Moore, A. R. (1). Some effects of temperature on development in the sea urchin Allocentrotus fragilis. Biol. Bull. Woods Hole 117: 150-153 4 figs.

Moore, A. R. (2). On the embryonic development of the sea urchin *Allocentrotus fragilis*. Biol. Bull. Woods Hole 117: 492-496 12 figs.

Moore, A. R. see Boolootian, R. A.

Morita, M. Histochemical determination of proteinbound sulfhydryl groups in the cortical layers of the sea urchin egg. Sci. Rep. Tôhoku Univ. 25: 45-51.

Morrison, J. F. see Griffiths, D. E.

Mosher, C. G. see Chaet, A. B.

Motomura, I. Secretion of mucosubstance in the cleaving eggs of the sea urchin. Bull. mar. biol. Sta. Asamushi 9 1958: 79.

Moura, A. Alguns equinideos regulares fósseis da Costa d'Arnes (Alfarelos, Coimbra). Mem. Mus. min. geol. Univ. Coimbra 45: 61-72 3 pls. figs. (Portuguese with French and English summaries).

Müller, H. see Zahn, R. K.

Mulnard, J. & others. Metachromasia observed in the living eggs of *Arbacia punctulata* and its cytochemical analysis. J. Embryol. exp. Morph. 7: 223-240 13 figs. 1 pl.

Nakamura, K. Acceleration of the cleavage in seaurchin eggs by heating prior to fertilization. Bull. mar. biol. Sta. Asamushi 9 1958: 63-64 1 fig.

Nakano, E. see Monroy, A.

Neefs, Y. Developpement et évolution sexuelle chez Asterina gibbosa. Proc. int. Congr. Zool. 15: 286-288.

Nichols, D. (1). Mode of life and taxonomy in irregular sea-urchins. Publ. Syst. Ass. No. 3: 61–80 7 figs.

Nichols, D. (2). Changes in the Chalk Heart-urchin *Micraster* interpreted in relation to living forms. Phil. Trans. 242B: 347-437 46 figs. pl. ix.

Nichols, D. (3). The histology of the tube-feet and clavulae of *Echinocardium cordatum*. Quart. J. micr. Sci. 100: 73-87 10 figs.

Nichols, D. (4). The history and activities of the tubefeet of *Echinocyamus pusillus*. Quart. J. micr. Sci. 100: 539-558 figs.

Nicola, M. G. de (1). I carotenoidi nell' intigumento degli Asteroidé: Astropecten aurantiacus. Ric. sci. 28 1958 : 2076–2083.

Nicola, M. G. de (2). I carotenoidi negli Asteroidi: Marthasterias glacialis. Ric. sci. 29: 78-84 1 fig.

Niewoehner, W. see Koenig, J. W.

Okada, M. see Tahara, Y.

Okazaki, K. Skeletal matrix in sea urchin lørvae. Bull. mar. biol. Sta. Asamushi 9 1958: 73.

Palinesar, E. E. Effects of some purine analogs on cleavage in *Arbacia punctulata*. Biol. Bull. Woods Hole 117: 436.

Parenzan, P. Biocenologia bentonica: il fondo ad Ascidie. Thalassia jonica 2: 15-45 4 figs.

Parker, R. H. Macro-invertebrate assemblages of central Texas coastal bays and Laguna Madre. Bull. Amer. Ass. Petrol. Geol. 43: 2100-2166 6 pls. 32 figs.

Perlmann, P. (1). Immunochemical analysis of the surface of the sea-urchin egg—an approach to the study of fertilization. Experientia 15: 41-52 17 figs.

Perlmann, P. (2). Immunochemical study of the macromolecules in the surface of the sea urchin egg. Proc. int. Congr. Zool. 15: 729-730.

Perlmann, P., Boström, H. & Vestermark, A. Sialic acids in the gametes of the sea urchin. Exper. Cell Res. 17 439-446 1 fig.

Poliansky, G. I. & Golikova, M. N. On the Infusoria dwelling in the intestine of sea urchins. III. Infusoria of sea urchins of the Barents Sea. Zool. Zh. 38: 1138-1145

[Poretskaya, E. S.] [The stratigraphical and paleogeographical distribution of the sea-urchins in the lower Tertiary deposits of the Mediterranean region.] Vesta. leningr. Univ. No. 24: 76-92 [Russian with English summary.]

Pressoir, L. Contribution à la connaissance des Échinoplutéus de Paracentrotus lividus Lmck., et Psammechinus microtuberculatus Blainv. océanogr. Monaco No. 1142: 1–22 16 figs.

Prosser, C. L. Activation of a non-propagating muscle in *Thyone*. J. cell. comp. Physiol. 44 1954: 247-253 5 for.

Rae, B. B. The food and feeding habits of the lemon sole. Mar. Res. 1956 No. 3: 1-32 5 figs.

Ramovš, A. Die Entwicklung des Oberperms in Loka- und Polhovgradechügel (W. Slowenien, NW. Jugoslawien). Razpr. Akad. Ljubljana 4 1958: 451–622 10 pls. 2 figs. 4 maps. [In Slovene with German summary on pp. 584–617].

Ranzi, S. (1). On the stability of the structures of animalized and vegetalized sea urchin embryos. Biol. Bull. Woods Hole 117: 436-437.

Ranzi, S. (2). The behaviour of antigens during the development of sea urchins. Proc. int. Congr. Zool. 15: 623-625.

Raup, D. M. Crystallography of Echinoid Calcite. J. Geol. 67: 661-674 7 figs.

Ray, C. Rococo creature of the sea—the basket starfish. Anim. Kingd. 62: 114-117 4 figs.

Reys, J. P. Thyone cherbonnieri nov. sp. et remarques sur le genre Thyone en Mediterranée. Rec. Trav. Sta. mar. Endoume No. 29: 173-180 6 figs.

Rocca, E. see Scarano, E.

Rodriguez, G. The marine communities of Margarita Island, Venezuela. Bull. mar. Sci. Gulf & Caribbean 9: 237–280 26 figs.

Roeder, M. The response of sea urchin plutei to added amounts of amino acid. Anat. Rec. 132: 495–496.

Rose, M. [in] Tregouboff, G. & Rose, M. Manuel de planctonologie méditerranéene. Chapter 33. Les larves du plancton. Paris 1957 vol. I pp. 485-510 vol. II pls. 157-191.

Roslansky, J. D. & Allen, R. D. The detection of rapid physical changes during fertilization of sea urchin eggs by the rate of elastic return following deformation. Anat. Rec. 132: 498.

Rothstein, H. The effect of sulfhydryl compounds on the colloidal state of the cytoplasm of *Arbacia* eggs. Biol. Bull. Woods Hole 117: 425.

Rulon, O. A comparison of the effects of nickelous chloride on the development of the sand dollar and the sea star. Anat. Rec. 132: 500.

Runnström, J. (1). Partial cortical activation of the sea urchin egg (*Paracentrotus lividus*). Ark. Zool. 12: 245-290 55 figs.

Runnström, J. (2). Inhibitory and anti-inhibitory effects on the activity of the centrosome in the sea urchin egg. Exper. Cell Res. 16: 448-451 1 fig.

Runnström, J. see Hagström, B. E.

Runnström, J. see Immers, J.

Rustad, R. C. (1). Consequences of unilateral ultraviolet radiation of sea urchin eggs. Biol. Bull. Woods Hole 116: 294-303 18 figs.

Rustad, R. C. (2). Further observations relating to radiation-induced mitotic delay to centriole damage. Biol. Bull. Woods Hole 117: 437.

Rustad, R. C. (3). The inhibition of mitosis in the sea urchin egg by acridine orange. Biol. Bull. Woods Hole 117: 437-438.

Rustad, R. C. (4). An interference microscopical and cytochemical analysis of local mass changes in the mitotic apparatus during mitosis. Exper. Cell Res. 16: 575–583 2 figs.

Saint-Seine, R. de. Proposition d'une nomenclature morphologique pour les aires coronales d'Echinides. Proc. int. Congr. Zool. 15: 1031-1032 1 fig.

Sakai, H. A fibrous protein and its sulfhydryl groups in dividing sea urchin eggs. Bull. mar. biol. Sta. Asamushi 9 1958: 65.

Sakai, H. & Dan, K. Studies on sulfhydryl groups during cell division of sea urchin egg. I. Glutathione. Exper. Cell Res. 16: 24-41 4 figs.

Sawada, N. Maturation division of clam eggs induced by the body fluids of sea urchins. Mem. Ehime Univ. (2) 3 1958: 1-3.

Scarano, E. & Rocca, E. Phosphoribose-isomerase from sea urchin sperm. Pubbl. Staz. zool. Napoli 31: 209–224 6 figs.

Scheffer, V. B. Invertebrates and fishes collected in the Aleutians 1936–38. N. Amer. Fauna No. 61: 365–406 26 figs.

Sebastio, C. Sugli Echinodermi eduli dei mari italiani e determinazione dei loro caratteri di vitalità e freschezza. Thalassia jonica 2: 82–108 10 figs.

Shao, T. see Dubatolowa, Y. A.

Sinclair, A. N. Observations on the behaviour of sea urchins. Aust. Mus. Mag. 13: 3-8 4 figs.

Souquet, P. Extension du Cénomanien au Sud du Massif de Castillon (Ariège). Bull. Soc. Hist. nat. Toulouse 94: 209-212 1 pl.

Speden, I. G. Phyllocrinus furcillatus sp. nov., a cyrtocrinoid from the Upper Jurassic of Kawhia, New Zealand. Palaeontology 2 1: 150–155 1 pl. 1 fig.

Speidel, C. C. & Cheney, R. H. Relation of ultravioletray dosage to modifications in fertilization and early development of *Arbacia*. Biol. Bull. Woods Hole 117 427.

Speidel, C. C. see Cheney, R. H.

Spiegel, M. & Spiegel, E. S. The extractable proteins of sea urchin embryos. Biol. Bull. Woods Hole 117: 427.

Stramondo, A. Il Neocomiano di Monte Boschitello presso Licodia Eubea (prov. Catania). Boll. Accad. gioenia 4 1958: 175–183.

Streble, H. (1). Die Entwicklung der Seeigel. Mikrokosmos 48: 176–181 12 figs.

Streble, H. (2). Zum Umschlagbild: Die Metamorphose der Seesterne. Mikrokosmos 48: 349-350 1 fig.

Suarez-Caabro, J. A. Salinidad, temperatura y plancton de las aguas costeras de Isla de Pinos. Monogr. Lab. Biol. mar., Habana Monogr. 7: 1–25 7 figs.

Sugiyama, M. The effect of nicotine on the sea urchin egg, with special reference to the fertilization-wave. Bull. mar. biol. Sta. Asamushi 9 1958: 53-54.

Sutherland, M. D. & Wells, J. W. Anthraquinone pigments from the crinoid Comatula pectinata. Chem. & Ind., 291-292.

Swami, K. S. see Wilson, W. L.

Szörényi, E. Les Torynocrinus (Crinoïdes) du Crétacé Inférieur de la Hongrie. Acta geol. hung. 6: 231-271 4 pls.

Tahara, Y., Okada, M. & Kobayashi, N. Secondary sexual characters in Japanese sea-urchins. Publ. Seto mar. biol. Lab. 7 1958: 165-172 2 pls. 5 figs.

Tambs-Lyche, H. Zoogeographical and faunistic studies on West Norwegian marine animals. Arb. Univ. Bergen. Naturv. R. 1958 7: 1-24 4 figs.

Termier, H. & Termier, G. see Cabibel, J.

Thieuloy, J.-P. Étude micrographique des "calcaires à débris" barrémo-aptiens sur le pourtour méridional du Vercors. Trav. Lab. Géol. Univ. Grenoble 35: 39-99 25 pls.

Tomita, H. see Aketa, K.

Tommasi, L. R. Os equinodermas do litoral de São Paulo. II. Diadematidae, Schizasteridae, Brissidae, Cidaroidae (Echinoidea) e Asteroidea do bento costeiro. Contr. avuls. Inst. oceanogr. S. Paulo No. 2 1958: 1-27 6 pls.

Tortonese, E. (1). Campagne de la Calypso dans le Golfe da Gênes (1957). 2. Échinodermes. Ann. Inst. océanogr. Paris N.S. 37 4 : 289–294.

Tortonese, E. (2). Ecofenotipi e biologia di Ophiothriz fragilis (Ab.) nel Golfo di Genova (1). Doriana 2 (100): 1-9.

Tweedell, K. S. Induced fluorescence in marine eggs. Biol. Bull. Woods Hole 117: 429-430.

Tyler, A. see Black, R. E.

Ubisch, L. von. Die Entwicklung von Genocidaris maculata und Sphaerechinus granularis, sowie Bastarde und Merogone von Genocidaris. Pubbl. Staz. Zool. Napoli 31: 159–208 45 figs. 3.pls.

Ushakov, B. Thermostability of the tissue as one of the diagnostic characters in poikilothermic animals. Proc. int. Congr. Zool. 15: 1046–1051 3 figs.

Utinomi, H. Coloured illustrations of sea-shore animals of Japan. Hoikushi Osaka 1956 pp. [2] xvii, 167: 64 col. pls., text figs. [Fauna and flora of Japan, No. 8]. [1. Japanese. Echinodermata on pp. 105–122 pls. liii–lxi.]

Vestermark, A. see Perlmann, P.

Vincent, W. S. & Baltus, E. The incorporation of radioactive label into RNA: synthesis or terminal addition. Biol. Bull. Woods Hole 117: 388.

Volk, M. Über fossilführende Gerölle im Ober-Ordovizium von Steinach, Thüringer Wald. Senck. leth. $\bf 40:$ 25–28 2 figs.

Wells, J. W. see Sutherland, M. D.

Went, H. A. (1). Some immunochemical studies on the mitotic apparatus of the sea urchin. J. Biophys. & Biochem. Cytol. 5: 353-356 1 fig.

Went, H. A. (2). Studies on the mitotic apparatus of the sea urchin by means of antigen-antibody reactions in agar. J. Biophys. & Biochem. Cytol. 6:447-455 4 pls.

Whiteley, A. H. see Litchfield, J. B.

Wilson, W. L. & Swami, K. S. Electrophoretic studies on protoplasm. Biol. Bull. Woods Hole 117: 389.

Wind, J. Echinocorys formerne og deres stratigrafiske Udbredelse i det øverste Kridt i Danmark. Medd. dansk geol. Foren. 14: 122–132 4 pls. [In Danish].

Yanagisawa, T. (1). Studies on echinoderm phosphagens. I. Occurrence and nature of phosphagens in seaurchin eggs and spermatozoa. J. Fac. Sci. Tokyo Univ. (Zool.) 8: 473–479.

Yanagisawa, T. (2). Studies on guanidine phosphoryltransferases of echinoderms. I. Occurrence in spermatozoa and eggs of sea-urchins. J. Fac. Sci. Tokyo Univ. (Zool.) 8: 481–486.

Yanagisawa, T. (3). Studies on the guanidine phosphoryltransferases of echinoderms. II. Some properties of the guanidine phosphoryltransferases of sea-urchin eggs and spermatozoa. J. Fac. Sci. Tokyo Univ. (Zool.) 8: 487–498 7 figs.

Yasumasu, I. (1). Hatching enzyme of the sea urchins Hemicentrotus pulcherrimus and Heliocidaris crassispina. Bull. mar. biol. Sta. Asamushi 9 1958: 83-84.

Yasumasu, I. (2). Changes in the content of echinochrome during embryonic development of the sea urchin, Hemicentrotus pulcherrimus. J. Fac. Sci. Tokyo Univ. (Zool.) 8: 499-503 3 figs.

Yoshida, M. Naphthaquinone pigments in *Psammechinus miliaris* (Gmelin). J. mar. biol. Ass. U.K. 38: 455–460 3 figs.

Yoshida, M & Millott, N. Light sensitive nerve in an Echinoid. Experientia 15: 13-14 3 figs.

Yoshida, M. see Millott, N.

Zahn, R. K. & Müller, H. Untersuchungen über die Zusammensetzung der Desoxyribonucleinsäure aus den reifen Spermatozoen von Seegurken. Z. Naturf. 13B 1958: 404-407.

Żakowa, H. [Biostratigraphy of the Lower Carboniferous marine deposits of the area of Walbrzych Miasto (Lower Silesia).] Trav. Serv. géol. Pologne 19 1958: 1–211 14 pls. 16 figs. [In Polish with English summary on pp. 183-205].

II.—SUBJECT INDEX

GENERAL LITERATURE AND HISTORY

Terminology.—Notation for coronal plates of echinoids, Saint-Seine.

Taxonomic characters.—Systematic serology of seastars, Kubo; irregular sea-urchins, Nichols (1) & (2); thermostability as a diagnostic character, Ushakov.

Collections.—Of Instituto Geologico y Minero de España, Badillo; echinoderms of Rumphius, Engel (with photographs of original drawings); Linnéan and Müllerian types in Copenhagen Zoological Museum, Madsen (2).

Techniques.—Statistical analysis of holothurian sclerites, Hampton.

STRUCTURE

General.—Anatomy of echinoid, Evechinus, McRae.

Skeleton.—Apical system of irregular echinoids, Jesionek-Szymanska; larval skeleton, Okazaki; crystallography of echinoid calcite, Raup.

Digestive system.—Cardiac stomach of Patiria, Anderson (2).

Tube feet.—Of Echinocardium, Nichols (3); of Echinocyamus pusillus, Nichols (4).

PHYSIOLOGY

Digestion.-Of starfish Patiria, Anderson (2).

Relation to environment.—Thermostability of tissues in poikilothermic animals, Ushakov.

Respiration.—Of unfertilized eggs, Brookbank; respiratory surface in an echinoid, Farmanfarmaian; oxygen equilibrium of haemoglobin in Cucumaria, Manwell; intensity of respiration in muscle. Mattigon.

Reactions to light.—Responses to increases in light intensity in echinoid Diadema antillarum, Millott & Yoshida; light sensitivity in an echinoid, Yoshida & Millott.

Pigments.—Carotenoids in Astropecten and in Marthasterius, Nicola (1) & (2); anthraquinone pigments from Comatula, Sutherland & Wells; naphthaquinone pigments in Psammechinus, Yoshida.

Regeneration.—Of pyloric caeca in starfishes, Anderson (1).

Nervous system.—Light sensitive nerve in an echinoid, Yoshida & Millott.

Metabolism .- Of Asterias, Bismarck.

Myoneural physiology.—Activation of longitudinal muscle of Thyone, Prosser.

Experimental studies.—Action of glycerol on protoplasm, Heilbrunn.

Bio-chemistry.—Effect of starfish toxin on amebocytes, Bang & Chaet; clotting of coelomic fluid, Boolootian & Giese; toxins from scalded starfishes, Chaet & Mosher; alginase in gut of echinoid, Eppley & Lasker; organic constituents of gonads, Giese & others; guanidines, phosphagens and N-amidino phosphokinases in echinoids, Griffiths, Morrison & Ennor; muscular extract of seacucumber, (In Japanese), Ishibashi; toxic effect of holothurin on frog, Jakowska & others; cytochrome o content and cytochrome oxidase activity in muscle, Mattison; echinoid body fluids inducing maturation division of clem eggs, Sawada; immunochemical studies of mitotic apparatus of sea-urchin, Went (1).

REPRODUCTION AND SEX

Reproductive cycles.—Seasonal gonad enlargement in Asterina gibbosa at Dinard, Delavault; increase in organic constituents of maturing gonads, Giese & others; development and sexual evolution in Asterina, Necis; spawning period of Strongylocentrotus pallidus, Tambs-Lyche.

Sex.—Sexuality of some mediterranean starfishes, Delayault & Cognetti.

Secondary sex characters.—In Japanese echinoids, Tahara, Okada & Kobayashi.

Hermaphroditism.—Natural ovotestis in echinoid, Boolootian & Moore.

DEVELOPMENT

Embryology.—Chromatin in pronucleus and syncarion of egg, Agrell (1); adhesion mechanism of cells in blastula and gastrula, Balinsky; presumptive position of dorsal and ventral areas in echinoid egg studied by local vitalstaining, Fukushi (1) & (2); cell mass on left side of Temnopleurus pluteus, Fukushi (3); factors influencing metamorphosis, Lynch; annulate lamellae in sand dollar eggs, Merriam; embryonic development of echinoid, Allocentrotus, Moore (2); skeletal matrix in larvae, Okazaki; larva of Paracentrotus lividus and Paammechinus microtuberculatus, Pressoir; larvae of Mediterranean Echinoderms, Rose; of Echinoids, Streble (1); metamorphosis of starfish, Streble (2); development of Genocidaris maculata from egg to metamorphosis and of Sphaerechinus granularis from young pluteus to metamorphosis, Ubisch.

Spermatosoa.—Sperm. tail, Afzelius; modifications of sperm. structure, Bernstein; influence of jelly coat solution on sperm., Hagström (1); inhibition of undulating in sperm., Kinoshits; phospholipid utilization during aging of sperm., Mohri; maturity of sperm., Zahn & Müller.

Fertilization including experimental studies.-Influence of temperature on nuclear charges during syngamy, Agrell (2); changes in cytoplasmic structure following fertilization, Allen & Bunim; inhibition of fertilization and agglutination by Fucus extract, Branham & Metz; effects of x-rays on fertilization, Chency & Speidel; membrane elevation in eggs, Costello; effect of temperature on fertilization, Hagström & Hagström; refertilization of partially fertilized eggs, Hagström & Runnström; effects of sperm.on S²⁶-labelled fertilizin, Hathaway; membrane elevation, Heilbrunn & Byers; changes in electric properties at fertilization, Hiramoto (2); change in membrane potential during fertilization of narcotized egg, Hori; toughening of fertilization membrane, Ishikawa; inhibiting effect of dermal secretion on fertilization, Metz (1) & (2); immunochemical approach to study of egg surface and fertilization, Perlmann (1) & (2); physical changes during fertilization detected by elasticity after deformation, Roslansky & Allen; partial cortical activation of echinoid egg, Runnström (1); modified activity of centrosome, Runnström (2); effect of ultraviolet-rays on fertilization, Speidel & Cheney.

Cleavage including cortical studies and mitosis.—Inhibition of cleavage and development by 2DG, Bernstein & Black; block to cell division produced by D₂O, Gross & others; effects of Na-salicylate on cleavage, Kitazume; factors in cleavage induction, Marsland; factors regulating cell division rate, Menkin; effects of substances from one species regulating cell division of another, Menkin & others; secretion of mucosubstance in cleavage, Motomura; acceleration of cleavage by heating before fertilization, Nakamura; effects of some purine analogs on cleavage, Palinesar; rhythmic changes in cytoplasm during cleavage cycles, Kojima; local mass changes in mitotic apparatus during mitosis, Rustad (1); inhibition of mitosis, Rustad (2) & (3); effect of nicotine on egg cortex, Sugiyama; mitotic apparatus studied by immunochemical methods, Went (1) and by antigen-antibody reactions in agar (2).

Experimental embryology.—Oxidative phosphorylation by unfertilized eggs, Aketa & Tomita; biological effect of copper in sea water tested by rate of echinoid development, Bougis; effects of metachromatic colorants on determination of eggs, Brice (2); toxicity of meta-

chromatic colorants for starfish eggs, Brice (4); respiration of unfertilized eggs in the presence of antisera against fertilizin, Brookbank (1); haptens of unfertilized eggs, Brookbank (2); negative redox potential in exogastrulation, Brooks; induction of gamete-shedding by nerve extracts, Chaet & McConnaughy; experiments on jellyree eggs, Hagström (2); hybridization experiments in echinoids, Hagström (3); electric properties of eggs, Hiramoto (1); effect of cyanide on recently fertilized eggs, Immers & Runström; effects of phenazone on embryonic determination, Lallier (1); SH groups in determination, Lallier (2); animalisation by zinc ions, Lallier (3); effect of Phenazone, Lallier (4); effect of ribonuclease on development, Leone; variations in thermal sensitivity during early development, Lenning; effects of temperature on echinoid development, Moore (1); effects of antisera on eggs, Perlmann (2); stability of structures in animalized and vegetalized embryos, Ranzi (1); behaviour of antigens in development, Ranzi (2); response of plutei to added amino acid, Roeder; effect of SH-compounds on colloidal state of cytoplasm, Rothstein: effects of nickelous chloride on development, Rulon; effect of unilateral ultraviolet radiation on eggs, Rustad (1); induction of maturation division in clam eggs by echinoid body fluid, Sawada; effect of ultravioletrays, Speidel & Cheney; induced fluorescence in eggs, Tweedell; development of hybrids between Genocidaris 2 and Paracentrotus or Psammechinus of and of hybrid merogones, Ubisch.

Chemical embryology.—Fucose in eggs, Ando; changes in RNA content during early development, Bāckström (1); SH-substances in embryos, Bāckström (2); oxidation of CO in echinoid eggs, Black & Tyler (1) & (2); oxidation of CO in echinoid eggs, Black & Tyler (1) & (2); oxidation of CO in echinoid eggs, Black & Tyler (1) & (2); oxidation skeletal formation of plutei, Hsiac; sea urchin egg proteins, Kane & Hersh; antigens of sperm.-surface, Kohler & Mets (1); extraction of jelly precipitating agents from sperm, Kohler & Mets (2); phosphate uptake by embryos, Litchfield & Whiteley; cytochrome oxidase in mitochondria, Maggic; analysis of lipids and lipoproteins of eggs and embryos, Marsh; localization of dehydrogenases, Merson; protein-bound SH groups in echinoid eggs, Morita; Metachromasia in living Arbacia eggs, Mulnard & others; immunochemical analysis of egg surface, Perlmann (1); sialic acids in gametes, Perlmann, Boström & Vestermark; fluctuation in -SH of fibrous protein during cell division, Sakai & Dan; phosphoribose-isomerase from echinoid sperm., Scarano & Rocca; extractable proteins of embryos, Spiegel & Spiegel; phosphagens in eggs and sperm., Yanagisawa (1), guanidine phosphoryl transferases, (2) and (3); hatching enzyme, Yasumasu (1); changes in echinochrome content during embryonic development, Yasumasu (2); DNA of Holothuria sperm., Zahn & Müller.

Techniques.—Electron microscopy of sperm., Afzelius; use of sea-urchin eggs in laboratory, Alcala; study of adhesion mechanism of cells by electron microscopy, Balinsky; study of skeletons of larvae with paired polarizing filters, Boolootian; method for measuring total lipid, Marsh; electron microscopy studies of sand dollar eggs, Merriman; methods for incorporation of radioactive compounds in echinoderm eggs, Monroy & Nakano; interference microscope in study of mitotic changes, Rustad (4); RNA labelling, Vincent & Baltus; immunochemical methods in study of mitotic apparatus, Went (1); electrophoretic studies on protoplasm, Wilson & Swami.

EVOLUTION AND GENETICS

Evolution.—Of the Microster group of fossils, Nichols (1) & (2).

Variation.—Of holothurian sclerites, Hampton; polymorphism of Ophiothrix fragilis, Tortonese (2).

Biometrics.-Of Micraster, Nichols (2).

ECOLOGY AND HABITS

Feeding habits.—Of starfish Patiria, Anderson (2); of Allocentrotus, Boolootian & others; of starfish Pisaster, Feder; of Echinus esculentus, Forster; Solaster, Asterias and Psammechinus in relation to each other and to oysters, Hancook; predators of mussels, Kitching & others; feeding behaviour in heart-urchins, Nichols (2); of basket-star in aquarium, Ray.

Behaviour.—Of echinoid Evechinus, McRae; response to increased light intensity in Diadema, Millott & Yoshida; of heart-urchins, Nichols (1) & (2); burrowing habits and 'homing' of sea-urchins, Sinclair.

Communities.—Animals associated with Allocentrotus, Boolootian & others; associated with Ophiothrix in the northern Adriatic, Czihak; benthic communities of Cook Strait, Hurley; bottom fauna of inshore fishing grounds off E. Scotland, McIntyre; of Ascidian beds in the Bay of Naples, Parenzan; of Texas coastal bays, Parker; of a Venezuelan island shore, Rodriguez.

Habitat.—Of deep-water Allocentrotus fragilis, Boolootian & others; of Odontaster validus in Ross Sea shown by photographs, Bullivant; of Ophiothrix in the northern Adriatio, Czihak; of echinoid Evechinus, McRae.

Environment.—Effects of salinity and temperature on plankton, Suarez-Caabro.

Distribution.—Quantitative distribution of Echinus esculentus. Forster.

Economics.—Old methods of cooking echinoderms in Amboina, Engel from Rumphius; poisonous echinoids, Halstoad; starfishes and Paammechinus as enemies of oysters, Hancock; food of lemon soles, Rae; edible echinoderms of Italian waters, Sebastio.

Techniques.—Underwater photography, Bullivant,

Parasites and Commensals.—Bryozoa epizoic on Leptometra, Gautier; rhabdocoel turbellarian in gut of Strongylocentrotus franciscanus, Giese; Dendrogaster tasmaniensis, crustacean parasite in starfish Allostichaster polyplax, Hickman; ascothoracid parasite in starfish Nepanthia belcheri, Kenny; Protozoan, rhabdocoel and copepod commensal with Evechinus chloroticus, McRae; Infusoria in intestines of Strongylocentrotus droebachiensis, Poliansky & Golikova.

DISTRIBUTION

GEOGRAPHICAL

Arctic.-Denmark Strait asteroid, Madsen (1).

North Atlantic.—Eastern U.S., echinoids, Cooke; Atlantic coast of France, Crisp & Fischer-Piette; Coney Island, New York, echinoids, Jakowska; Brittany, Paracentrotus, Lami; Dorset, England, Asterias rubens, Lang; East of Scotland, McIntyre; W. Norway, Tambs-Lyche.

Tropical Atlantic.—Brazil, Brito; French Guiana, Cherbonnier (1); Florida, holothurians, Domantay (2);

Gulf of Mexico, Parker; Venezuela, Rodriguez; Brazil, echinoids and asteroids. Tommasi.

Mediterranean.—Thyone, Reys; key to echinoderm larvae of the Mediterranean, Rose; Italy, Tortonese (1).

Indo-W. Pacific.—Red Sea Holothurians, Cherbonnier (2); Fauna list of Holothurians, Domantay (1); New Zealand Asteroids, with key to spp., Fell; Great Barrier Reef, Gillett & McNeill; ophiuroids in Cook Strait, New Zealand, Hurley; New Zealand, echinoid, McRae; Japan, Iffinowi

East Pacific.—The Ekman barrier and Panamanian holothurians, Deichmann (1); California, with key to intertidal species. Light & others.

North Pacific.-Aleutian Is., Scheffer.

South Temperate.—S. Brazil, holothurians, Ancona-Lopez; S. Atlantic coasts of Africa, echinoids, Cherbonnier (3).

Antarctic.—Holothurian fauna, Deichmann (2); asteroid, Fell & Clark.

GEOLOGICAL.

Holocene.-Angola, Andrade & Andrade.

Pleistocene.—Angola, Andrade & Andrade; eastern U.S., echinoids, Cooke.

Pliocene.—Angola, Andrade & Andrade; eastern U.S., echinoids, Cooke.

Miocene.—Angola, Andrade & Andrade; eastern U.S., echinoids, Cooke; Trinidad, Jeannet.

Oligocene.—Angola, Andrade & Andrade; eastern U.S., echinoids, Cooke; Holstein, N. Germany, holothurians, Deflandre-Rigaud; Trinidad, Jeannet; Mediterranean region, especially southern U.S.S.R, Poretskaya.

Eccene.—Angola, Andrade & Andrade; France, Balavoine; eastern U.S., echinoids, Cooke; Trinidad, Jeannet; Mediterranean region, especially southern U.S.S.R., Poretskaya.

Paleocene.-Eastern U.S., echinoids, Cooke.

Cretaceous.—Angola, Andrade & Andrade; Texas, U.S., ophiuroids, Clark, D. L.; Angola, Ferreira Soares (1) & (2); Georgia, USSR, Kotetishvili; Austria, Leischner; E. Serbia, Mitrović-Petrović; Portugal, echinoids, Moura; Pyrenees, Souquet; Sicily, Stramondo; Hungary, crinoids, Szörenyi; S.E. France, Thieuloy; Denmark, Wind.

Jurassic.—Irregular echinoids, Jesionek-Szymanska; Austria, Leischner; New Zealand, crinoid, Speden.

Triassic.—Southern China, crinoids, Dubatolowa & Shao.

Permian.—Southern China, crinoids, Dubatolowa & Shao; Kansas, U.S., ophiuroid, Hattin; W. Slovenia and N.W. Jugoslavia, Ramovš.

Carboniferous.—Southern China, crinoids, Dubatolowa & Shao; Indiana, U.S., holothurians, Gutschick; Lower Silesia, Poland, Zakowa.

Devonian.—Pennsylvania, U.S., starfish, Cramer; Missouri, U.S., Koenig & Niewoehner; N. Africa, Le Matre.

Ordovician.-Germany, Volk.

Cambrian.—Esp. Spain, Badillo; S. France, Cabibel, Termier & Termier.

III.—SYSTEMATIC INDEX

HOLOTHURIOIDEA

†Achistrum brevis sp. nov. Carboniferous Indiana, U.S. pp. 132-133 text-fig. 2 pl. xxv figs. 20-24, ludwigi p. 133 pl. xxv figs. 13-19, triassicum? pp. 133-134 pl. xxv figs. 25, 26, Gutschick.

Actinopygu agassizi, Domantay (2) p. 190, with detailed synonymy.

Afrocucumis africana, Utinomi p. 122 pl. lxi fig. 8.

†Calclamnella p. 192 with C. fusiformis sp. nov. Oligocene Holstein, Germany p. 192 pl. i figs. 8-9 pl. ii fig. 11, Deflandre-Rigaud.

†Calcligula p. 192 with C. elgeri sp. nov. Oligocene Holstein, Germany, Deflandre-Rigaud pp. 192-193 pl. i figs. 14-17 pl. ii fig. 4 pl. iii figs. 1-4.

Chiridota rotifera, Ancona-Lopez pp. 17-20 figs. 45-50.

Cucumaria saxicola, Hampton p. 344 text-fig. 2 pl. i fig. 3 pl. iv.

†Cucumarites p. 191 with C. holsaticus p. 191 pl. i figs. 1, 2 pl. ii figs. 1-3, scaber pp. 191-192 pl. i fig. 3 pl. ii fig. 5, speciosus p. 192 pl. i fig. 4 pl. ii fig. 6, spp. nov. Oligocene Holstein, Germany, Dellandre-Rigaud.

†Dictyothurites gen. nov. super-order Actinopoda, for type D. corbisema sp. nov. p. 193 pl. i figs. 12, 13 pl. ii fig. 10, spatuligerus sp. nov. p. 193 pl. i figs. 10, 11 pl. ii fig. 13, both Oligocene Holstein, Germany, Deflandre-Rigand.

†Elgerius gen. nov. super-order Actinopoda, for type E. ostrea sp. nov. p. 192 pl. i fig. 5 pl. ii fig. 7 pl. iii fig. 14c, E. innienensis sp. nov. p. 192 pl. i fig. 6 pl. ii figs. 8, 9 pl. iii figs. 14a, b, both Oligocene Holstein, Germany, Deflandre-Rigaud.

Euapta godeffroyi, Domantay (2) pp. 198-199, with detailed synonymy.

Euthyonidiella zacae, Domantay (2) p. 192, with synonymy.

Holothuria (Holothuria) grisea, Ancona-Lopez pp. 7-11 figs. 1-24; H. arenicola pp. 183-184, atra p. 183, cubana pp. 184-185, floridana p. 185, glaberrima p. 186, grisea pp. 186-187, impatiens pp. 187-188, mexicana p. 188, surinamensis pp. 188-189, tubulosa pp. 189-190, with detailed synonymies, Domantay (2); H. leucospilota Gillett & MoNeill pl. cliii; H. impatiens, Hampton pp. 340-344 text-figs. 1, 3 pls. i figs. 1, 2 pls. ii, iii; H. leucospilota p. 121 pl. lxi fig. 5, lubrica var. moebi p. 121 pl. lxi fig. 6, monacaria pp. 120-121 pl. lxi fig. 3, pervicax p. 121 pl. lxi fig. 4, Utinomi.

Leptopentacta deichmannae pp. 193-194, nova var. floridana p. 194, nova var. gemmata pp. 194-195, Domantay (2).

Ludwigia americana, Domantay (2) p. 195.

†Microantyx botoni sp. nov. Carboniferous Indiana, U.S., Gutschick pp. 134–135 text-fig. 3 pl. xxvi figs. 22, 23, 28, 29.

†Myriotrochites p. 194 with M. intercessus pp. 196, 198 pl. i fig. 22 pl. iii fig. 11, mirabilis pp. 194, 196 pl. i figs. 18-20 pl. iii figs. 7-9, 12, undosus p. 196 pl. i fig. 21 pl. iii fig. 10, spp. nov. Oligocene Holstein, Germany, Deflandre-Rigaud.

Neothyone capensis var. trojani, Domantay (2) pp. 192-193.

Estergrenia digitata, Tortonese (1) p. 290.

Paracaudina chilensis var. ransonneti, Utinomi p. 122 pl. lxi fig. 9.

Pentacta australis, Utinomi p. 121 pl. lxi fig. 7.

Phyliophorus (Urodemella) occidentalis, Domantay (2) pp. 191-192, with detailed synonymy.

Polycheira rufescens, Utinomi p. 122 pl. lxi fig. 11.

Polyplectana kefersteinii, Domantay (2) p. 199, with detailed synonymy.

†Priscopedatus p. 193 with P. conspicuus sp. nov. Oligocene Holstein, Germany p. 193 pl. i fig. 7 pl. ii fig. 12 pl. iii fig. 6, Deflandre-Rigaud.

Protankyra bidentata, Utinomi p. 122 pl. lxi fig. 10.

†Rota campbelli sp. nov. Carboniferous Indiana, U.S., Gutschick pp. 135-136 text-fig. 4 pl. xxvi figs. 1-21, 26, 27, 32-35.

Stichopus badionotus, Ancona-Lopez pp. 11-14 figs. 25-33, 57, Cherbonnier (1) pp. 440-443 fig. 10a-m, fig. 11a-g, Domantay (2) pp. 190-191, with detailed synonymy; S. chloronotus p. 120 pl. lxi fig. 2, japonicus p. 120 pl. lxi fig. 1, oshimae p. 120 pl. lx fig. 6, Utinomi.

Synapta maculata, Gillett & McNeill p. 103 pl. xcix.

†Synaptites (Calcancora) spectabilis sp. nov., Deflandre-Rigaud p. 198 pl. i figs. 23, 24 pl. iii fig. 13.

Synaptula secreta sp. nov. S. Brazil, Ancona-Lopez pp. 20-26 figs. 51-56.

Thelenota ananas, Utinomi p. 120 pl. lx fig. 5.

†Thuroholia spicatus sp. nov. Carboniferous Indiana, U.S. p. 132 text-fig. 1 pl. xxv figs. 3, 4, 7-9, maryinata p. 131 pl. xxv figs. 2, 6, mecormacki p. 131 pl. xxv figs. 1, 5, Gutschick.

Thyonacta sabanillaensis, Cherbonnier (1) p. 440.

Thyone cherbonnieri sp. nov. S. France pp. 173-175 figs. 1-6, T. fusus, gadeana pp. 176, 177, Reys; T. (Sclerodactyla) braziliensis, Ancona-Lopes pp. 14-17 figs. 34-44; T. clarki pp. 195-196, floridana p. 196, fusus pp. 196-197, pervicax p. 197, pseudofusus pp. 197-198, tenella p. 198, Domantay (2).

Trochodota japonica, Utinomi p. 122 pl. lxi fig. 12.

New genus?, new species, **Gutschick** p. 136 text-fig. 5 pl. xxvi figs. 24, 25, 30, 31.

Beche-de-mer, Gillett & McNeill pl. xcvi figs. 1-3 (expelling viscera).

ECHINOIDEA

†Abertella p. 44, with A. aberti pp. 44-45 pl. xvi figs. 1-3, Cooke.

Acrosalenia angularis, Jesionek-Szymanska text-fig. 3A.

†Agassizia p. 74, with A. floridana p. 75 pl. xxxii figs. 1-4, porifera pp. 74-75 pl. xxxi figs. 1-8, A. (Anisaster) p. 75, A. (Anisaster) mossomi p. 76 pl. xxxii figs. 5-9, wilmingtonica p. 76 pl. xxxii figs. 10-13, Cooke.

Allocentrotus fragilis, Boolootian & others fig. 1 A-E.

 $\dagger Amblypygus$ pp. 26–27, with A. americanus p. 27 pl. vii figs. 8, 9, Cooke.

Anthocidaris crassispina, Utinomi pp. 115-116 pl. lviii fig. 5.

Arachnoides placenta, identity of 'Echinus planus' of Rumphius, 1705 p. 36, 1 pl. xiv fig. G, Engel p. 216 photo 24 fig. 3.

Arbacia aequituberculata, Sebastio pp. 100-101 figs. 7, 8; A. lizula var. africana, Cherbonnier (3) pp. 43 pl. iii figs. G-P pl. iv. fig. A; A. punctulata, Cherbonnier (1) p. 367, Cooke p. 19, Jakowska pp. 137-139 2 figs.

†Arbacia p. 19, with A. improcera pp. 20-21 pl. iv figs. 1-3, punctulata Pleistocene and Recent p. 19 pl. iv figs. 7-9, rivuli p. 20 pl. iv figs. 10-11, sloani p. 21 pl. iv figs. 12-14, vaccamav pp. 19-20 pl. iv figs. 15-17, Cooke.

†Arbia p. 21, with A. aldrichi pp. 21-22 pl. iii figs. 15-17, Cooke.

†Archaeocidaris urii, Zakowa p. 127 pl. x figs. 5a-d.

Asthenosoma ijimai, Halstead figs. 23, 24.

Astriclypeus manni, Utinomi p. 118 pl. lix fig. 9.

Astropyga magnifica, Cherbonnier (1) p. 368; A. nuptialis, Tommasi pp. 3-4 pl. i fig. 1.

Brissopsis caparti sp. nov. Gulf of Guinea pp. 51-54 pl. ix figs. A-O, atlantica var. mediterranea p. 51 pl. viii figs. M-O, jarlii pp. 54-56 text-fig. 1 pl. x figs. A-L, B. sp. pp. 56-57, Cherbonnier (3); B. atlantica, Cooke p. 85 pl. xxxviii figs. 19-22; B. lyrifera, Nichols (2) pp. 356, 376-377 fig. 5d-f, 8d-f, Tortonese (1) p. 292.

†Brissopsis p. 84, with B. biarritzensis p. 85 pl. xxxviii figs. 14–18, bianpiedi pp. 85–86 pl. xxxviii figs. 1–8, steinhatchee p. 85 pl. xxxviii figs. 9–13, Cooke.

Brissus latecarinatus, Utinomi p. 119 pl. lx fig. 1; B. unicolor, Cooke pp. 81-82 pl. xxxvi figs. 1-4.

†Brissus p. 81, with B. glenni sp. nov. Miocene S. Carolina, U.S. p. 82 pl. xxxvi figs. 5, 6, unicolor pp. 81–82 pl. xxxvi figs. 1-4 Pleistocene to Recent, Cooke.

†Brochopleurus p. 18 with B. pretiosus pp. 18-19 pl. iv figs. 4-6, Cooke.

Cassidulidae, with key to genera and spp. from eastern U.S., Cooke p. 56.

†Cassidulus pp. 56-57, with C. ericsoni p. 58 pl. xxiv figs. 13-16, gouldii pp. 57-58 pl. xxiv figs. 5-12, sabistonensis p. 57 pl. xxiii figs. 6-14, trojanus pp. 58-59 pl. xxiv figs. 1-4, C. (Plagiopygus) p. 59, with C. (Plagiopygus) conradi pp. 59-60 pl. xxv figs. 3-6, conradi carolinensis p. 60 pl. xxv figs. 7-10, conradi lyelli p. 60 pl. xxv figs. 1, 2, georgianus pp. 60-61 pl. xxv figs. 14-17, georgianus globosus p. 61 pl. xxv figs. 11-13, Cooke.

Centrostephanus longispinus, Cherbonnier (3) pp. 44-45 pl. iv figs. B-L pl. v figs. A, B; C. rodgersii, Sinclair p. 5 fig.

Cidaris p. 8 with C. abyssicola p. 8, Affinis p. 9, tribuloides p. 9 pl. i figs. 1, 2, Cooke; C. cidaris var. meridionalis, Cherbonnier (3) pp. 39-41 pl. i figs. A-I pl. ii figs. A-E.

†Cidaris figueirensis, Moura pp. 62-63 text-fig. 1 pl. i figs. 1, 2; C. Malheiroi, Ferreira Soares (1) p. 19 pl. v fig. 8, (2) p. 149 pl. v fig. 8; C. pratti p. 10 pl. i figs. 6, 18, 19, splendens pp. 9-10 pl. i fig. 10, C. sp. aff. splendens p. 10 pl. i fig. 11, Cooke.

i

6

Clypeaster (Orthanthus) aloysioi sp. nov. Brazil, Brito pp. 1-4 pls. i, ii; C. durandi sp. nov. French Guiana, pp. 370-372 fig. 8a-e fig. 9a-d, subdepressus pp. 368-370, Cherbonnier (1); C. rosaccus p. 34 pl. x figs. 1-3, subdepressus p. 36 pl. xi figs. 2-4, Cooke; C. rosaccus, possible identity of Rumphius, 1705 pl. xiv fig. C, Engel p. 215 photo 24 fig. 4; Clypeaster japonicus p. 117 pl. lix fig. 3, reticulatus p. 117 pl. lix fig. 2, virescens p. 117 pl. lix fig. 1,

†Clypeaster pp. 33-34 with key to spp. from eastern U.S., C. concavus pp. 34-35 pl. x figs. 4-6 pl. xi fig. 5, cotteau pp. 37-38 pl. xii figs. 1-3, oxybaphon pp. 35-36 pl. xi fig. 1, rogersi pp. 36-37 pl. xii figs. 4-6, rosaceus Pleistocene and Recent p. 34 pl. x figs. 1-3, subdepressus Pleistocene and Recent p. 36 pl. xi figs. 2-4, Cooks.

†Coelopleurus p. 22, with C. infulatus pp. 22-23 pl. iii figs. 5, 6, Cooke.

†Collyrites sp., Jesionek-Szymanska pp. 341–342 fig.

Colobocentrotus atratus, Madsen (2) pp. 167–168 fig. 3, notes on the holotype; C. mertensi, Utinomi, p. 116 pl. lviii fig. 8.

†Cyclaster p. 68, with C. drewryensis pp. 68-69 pl. xxviii figs. 1-4, Cooks.

Diadema savignyi p. 113 pl. lvii fig. 5, setosum p. 113 pl. lvii fig. 4, Utinomi; D. setosum identifiable from description of Rumphius, 1705 p. 35 though pl. xiii fig. 5 is 'peculiar,' Engel p. 214 photo 23 fig. 2, pl. xiv fig. B 'Diadema turcarum' is also probably setosum p. 211 photo 23 fig. 11; D. setosa Gillett & McNeill pl. xciv, D. setosum, Halstead fig. 23, Tahara, Okada & Kobayashi p. 170 fig. 5 pl. viii figs. 9, 10.

†Diplopodia marticensis pp. 64-66 text-fig. 2 pl. i figs. 3, 4 pl. ii figs. 1-3, variolaris pp. 68-69 pl. ii fig. 4, Mours.

Disaster sp. nov., Jesionek-Szymanska p. 342 text-fig. 2 pl. i fig. 3.

† Ditremaster p. 76, with D. becker ip. 77 pl. xxx figs. 9-12, Cooke.

†Dixieus p. 23 with D. dixie p. 24 pl. v figs. 8-10, Cooke.

†Echanthus p. 62, with E. georgiensis pp. 62-63 pl. xxvi figs. 13-16, Cooke.

Echinarachnius p. 44, with E. parma p. 44 pl. xiii figs. 5, 6, Cooke.

Echinocardium cordatum, Cooke p. 78, Jesionek-Szymanska fig. 3B, C, Utinomi p. 119 pl. lx fig. 2; E. cordatum pp. 355, 368-373, 375-376 figs. 1d-f, 4d-f, 7d-f, 11, 13, 15 (part), 16-19, 26-30 pl. ix figs. 47-50; flavescens pp. 355, 376 fig. 3a-c, 6a-c, 9a-c; pennatifidum pp. 355, 376 fig. 3d-f, 6d-f, 9d-f, Nichols (2).

†Echinocardium p. 78, with E. gothicum p. 79 pl. xxxiii figs. 7-10, orthonotum pp. 78-79 pl. xxxiii figs. 1-5, Cooke.

†Echinoconus castanea, Mitrović-Petrović pp. 50-51 pl. ii fig. 2.

†Echinocorys pp. 122–123 with E. legindensis sp. nov. p. 125 pl. ii figs. 1–6, pustolosus daniensis subsp. nov. pp. 127–128, both Cretaceous Denmark, E. brevis p. 130, minor pp. 128–130 pl. iii figs. 7–9, pustolosus pustolosus pp. 126–127, scutatus p. 124 pl. i figs. 1–6, sulcatus sulcatus pp. 125–126, sulcatus forma a p. 126, E. sp. nov. (aff. sulcatus forma a) p. 126, Wind; Echinocorys p. 66, with E, ovalis pp. 66–67 pl. xxvii figs. 6–9, Cooke.

Echinocyamus pusillus, Nichols (2) pp. 356-358, 374, 377 fig. 10.

†Echinocyamus p. 31, with E. macneili sp. nov. Eocene Alabama, U.S. pp. 32-33 pl. ix figs. 6-8, chipolanus p. 32 pl. ix figs. 1-3, hualeyanus p. 32 pl. ix figs. 12-14, meridionalis p. 32 pl. ix figs. 4, 5, parvus p. 31 pl. ix figs. 9-11, Cooks. Echinodiscus auritus, identity of specimen with two open slits on p. 37 and pl. xiv fig. F of Rumphius, 1705, Engel p. 216 photo 24 fig. 8, also E. bisperforatus 'young ones.'

Echinolampas ovata, identity of Rumphius' pl. xiv fig. 3, 1705, Engel p. 216 photo 24 fig. 6.

†Echinolampas (Psammolampas) kugleri sp. nov. Oligocene Trinidad pp. 201-203 pl. ii figs. 4-6 pl. ix, E. (Cypholampas) lycopersicus pp. 196-197 pls. i, ii fig. 1-3, iii, iv, v and vi, E. cf. lycopersicus pp. 197-198 pl. vii figs. 1-3, E. lycopersicus var. p. 198 pl. vii figs. 4-6, Jeannet; Echinolampas p. 55, with E. aldrichi pp. 55-56 pl. xxii figs. 11-13, appendiculata p. 56 pl. xxii figs. 5, 6, Cooke.

Echinometra p. 25, with E. licunter p. 25 pl. vi figs. 1, 2, Cooke; 'Echinometra digitata prima loblonga' of Rumphius 1705, p. 32, I 'most probably' is Hetericentrotus mammillatus, Engel pp. 212-213, 'Echinometra digitata secunda l. notunda' Rumphius p. 33, II 'not identifiable with any degree of certainty' pp. 213-214, 'Echinometra setosa' Rumphius p. 35 is type of Diadema setosum (Leske), p. 214; 'Echinus saxatilis' of p. 31, II, is probably Echinometra mathaei p. 211; E. mathaei, Gillett & McNeill pl. xev fig. 3, Tahara, Otada & Kobayashi p. 169 pl. viii figs. 7, 8, Utinomi p. 116 pl. lviii fig. 7; E. oblonga, Alcala pl. ii.

Echinoneus cyclostomus, identity of Rumphius' 1705, pl. xiv fig. D, Engel p. 215 photo 24 fig. 7; E. cyclostomus with synonym Echius oralis Linnaeus 1753, Madsen (2) pp. 168-169 fig. 4 (? holotype of oralis); E. cyclostomus, Utinomi p. 118 pl. lix figs. 10, 11.

†Echinopsis p. 14 with E. diatreta p. 14 pl. ii fig. 8-11, Cooke.

Echinostrephus aciculatus, Tahara, Okada & Kobayashi, pp. 169-170 fig. 4, Utinomi p. 116 pl. viii fig. 6.

Echinothrix calamaris p. 114 pl. lvii fig. 7, diadema pp. 113-114 pl. lvii fig. 6, Utinomi.

Echinotrix diadema, probable identity of 'Echinus niger' of Rumphius 1705 p. 31, III, Engel p. 211.

Echinus acutus, Cherbonnier (3) pp. 46–48 pl. v figs. C-J pl. vi figs. A-F; Echinus esculentus of Rumphius, 1705, p. 31, I description of Tripneustes gratilla but pl. xiii figs. B, C correctly named, Engel pp. 210–211 photo 23 figs. 7, 9, E. niger of p. 31, III probably Echinotrix diadema p. 211, E. planus of p. 36, I and pl. xiv fig. G is Arachnoides placenta p. 216 photo 24 fig. 3, E. saxatilis of p. 31, II is probably Echinometra mathaei p. 211 but pl. xiv fig. A might be Diadema setosum photo 23, fig. 15, E. sulcatus of p. 36 unidentifiable but pl. xiv fig. C is presumably Clypeaster rosaceus p. 215 photo 24 fig. 4; E. melo, Sebastio p. 101 figs. 9, 10.

Encope emarginata, Cherbonnier (1) p. 372; E. emarginata, pp. 49-50 pl. xviii fig. 5 pl. xviii fig. 1, michelini p. 49 pl. xviii figs. 2, 3, Cooke.

†Encope pp. 47–48 with key to spp. from eastern U.S., E. emarginala Late Micoene (?) to Recent pp. 49–50 pl. xvii figs. 5 pl. xviii figs. 1, macrophora p. 48 pl. xvii figs. 1, 2, michelini Pleistocene and Recent p. 49 pl. xviii figs. 2, 3, tamiamiensis pp. 48–49 pl. xvii figs. 3, 4, Gooke.

Eucidaris tribuloides var. africana, Cherbonnier (3) pp. 41-43 pl. ii figs. F-O pl. iii figs. A-F; E. tribuloides, Tommasi pp. 4-5 pl. i fig. 2; E. metularia, Utinomi p. 113 pl. Ivii fig. 2.

†Eupatagus pp. 88-89, with E. clevei p. 89 pl. xli figs. 6-8, E. (Gymnopatagus) pp. 89-90, E. (Gymnopatagus) antillarum p. 90 pl. xli figs. 1-5, carolinensis p. 91 pl. xlii figs. 4-7, gardnerae pp. 90-91 pl. xlii figs. 1-3, ocalanus pp. 91-92 pl. xlii figs. 8-12, E. (Brissopatagus) p. 92, E. (Brissopatagus) alabamensis p. 92 pl. xliii figs. 5-11, Cooke.

†Eurhodia p. 63, with E. ? elbana pp. 64-65 pl. xxvi figs. 9-12, E. patelliformis p. 64 pl. xxii figs. 7-10, rugosa pp. 63-64 pl. xxii figs. 1-4, E. (Gisopygus) p. 65, E. (Gisopygus) holmesi p. 65 pl. xxiii figs. 1-5, Cooke.

Evechinus chloroticus, all aspects, McRae pp. 205-266 figs. 1-15 pls. xii, xiii.

Fibularia acuta, Utinomi p. 118 pl. lix fig. 12.

†Fibularia p. 30, with F. alabamensis sp. nov. Eocene Alabama, U.S. p. 31 pl. ix figs. 20-22, texana p. 30 pl. ix figs. 15-19, vaughani pp. 30-31 pl. ix figs. 23-27, Cooke.

†Gagaria p. 17 with G. chickasawhay p. 18 pl. iii fig. 18, mossomi pp. 17-18 pl. iii figs. 10-14, salis p. 17 pl. iii figs. 7-9, Cooke.

†Galerites vulgaris, possible identity of Rumphius', 1705, pl. 1 fig. 8 and ? also 6, 7 and 9, Engel p. 222.

†Gauthieria p. 23, with G. speciosa p. 23 pl. v figs. 1-7, Cooke; G. broecki, possible identity of Rumphius', 1705, pl. lix fig. E, Engel p. 222.

Heliocidaris erythrogramma, Sinclair p. 6 fig.

†Hemiaster p. 67, with H. moscovensis sp. nov. Paleocene Alabama, U.S. p. 68 pl. xxviii figs. 5–9, parastatus p. 67 pl. xxviii figs. 15–20, stella pp. 67–68 pl. xxviii figs. 10–14, Cooke.

Hemicentrotus pulcherrimus, Utinomi p. 115 pl. lviii fig. 3.

Heterocentrotus mammillatus, probable identity of 'Echinometra digitata prima loblonga' of Rumphius, 1705 p. 32, I, pl. xiii figs. 1, 2 and D, Engel pp. 212–213 photo 23 figs. 6, 16, 10, 12, 13; H. mamillatus, Gillett & McNeill pl. xcv fig. 1, Utinomi p. 116 pl. lviii fig. 9.

†Heterodiadema ouramense, Moura pp. 69-71 pl. iii figs. 1, 2.

†Holaster p. 65, with H. cinctus pp. 65-66 pl. xxvii figs. 1-5, Cooke; H. lerichei, Ferreira Soares (1) p. 20 pl. v figs. 5, 6, (2) pp. 150-151 pl. v figs. 5, 6; H. complanatus pp. 48-49 pl. i fig. 4, prestensis pp. 49-50 pl. ii fig. 1, Mitrović-Petrović.

Laganum laganum 'may well be 'the 'second species' of Rumphius, 1705, p. 37 pl. xiv fig. E, Engel pp. 216-217 photo 24 fig. 11.

†Laganum p. 50, with L. floridanum pp. 50-51 pl. xx figs. 8-10, ocalanum p. 51 pl. xx figs. 11-15, Cooke.

Leodia p. 46, with L. sexiesperforata pp. 46-47 pl. xix figs. 4, 5, Cooks.

†Leodia caroliniana, Cooke p. 47 pl. xix figs. 1-3.

†Linthia p. 69, with L. alabamensis pp. 69-70 pl. xxix figs. 5-7, hanoverensis p. 70 pl. xxix figs. 8-12, L. ? prima p. 70 pl. xliii figs. 1-4, wilmingtonensis p. 69 pl. xxix figs. 1-4, Cooke.

Lovenia elongata, Utinomi p. 119 pl. lx fig. 3.

†Lovenia p. 77, with L. alabamensis sp. nov. Oligocene Alabama, U.S. p. 77 pl. xxxii figs. 14–17, clarki pp. 77–78 pl. xxxiii fig. 6, Cooke.

Lytechinus variegatus, Cooke, p. 15 pl. ii figs. 12-13.

†Lytechinus pp. 14-15, with L. floralanus p. 15 pl. ii figs. 17-21, variegatus Pleistocene and Recent p. 15 pl. ii figs. 12-13, Cooke.

Macropneustes ventricosus, Cooke p. 84 pl. xxxvii figs.

†Macropneustes pp. 82-83, with M. mortoni pp. 83-84 pl. xxxvii figs. 5-9, Cooke.

†Maretia pp. 80-81, with M. arguta p. 81 pl. xxxiv figs. 1-4, subrostrata p. 81 pl. xxxiv figs. 5, 6, Cooke.

Mellita pp. 45-46, with M. quinquiesperforata p. 46 pl. xix figs. 6, 7, Pleistocene and Recent, quinquiesperforata tenuis p. 46, Cooke.

 $\dagger \textit{Mellita quinquies perforata}$, Cooke, p. 46 pl. xix figs. 6, 7.

Meoma Gray 1851 a synonym of Macropneustes Agassiz in Agassiz and Desor 1847, Cooke, p. 83.

Mespilia globulus, Tahara, Okada & Kobayashi pp. 166–168 figs. 1, 2 pl. vii figs. 1, 2, Utinomi p. 114 pl. lvii fig. 8.

Metalia spatagus, identity of Rumphius', 1705, pl. xiv, fig. 1, Engel p. 215 photo 24 fig. 10.

†Micraster, possible identity of Rumphius', 1705, pl. lix fig. F, Engel p. 223; Micraster pp. 401-427 with M. coranguinum figs. 40-43, 45, senonensis fig. 46, Nichols (2).

Moira p. 73, with M. atropos Pleistocene and Recent pp. 73-74 pl. xxx figs. 1-4, Cooke; M. atropos, Tommasi pp. 5-6 pl. i fig. 3.

†Moira atropos, Cooke pp. 73-74 pl. xxx figs. 1-4.

†Mortonella Pomel 1883 a synonym of Periarchus Conrad 1866, Cooke, pp. 41, 44.

†Neolaganum p. 51, with N. durhami sp. nov. Eocene Florida U.S. p. 52 pl. xxi figs. 5-7, dalli pp. 51-52 pl. xxi figs. 1-4, Cooke.

Neolampas rostellata, Tortonese (1) p. 292.

'Oculus Poliphemi' of Rumphius 1705 p. 31, IV, unidentifiable, **Engel** pp. 211, 212.

†Oligopygus pp. 27-28, with O. haldemani p. 29 pl. viii figs. 6-8, rotundus pp. 29-30 pl. viii figs. 1-5, wetherbyi p. 28 pl. viii figs. 9-12, Cooke.

†Orthopsis sp., Ferreira Soares (1) p. 20 pl. v fig. 7, (2) p. 150 pl. v fig. 7.

Paracentrotus gaimardi, Cherbonnier (3) pp. 48-49 pl. vi figs. G-K pl. vii figs. A-J; P. lividus, Crisp & Fischer-Piette pp. 327-328, Sebastio p. 99 figs. 1-4, Tortonese (1) p. 292.

†Paraster p. 71, with P. americanus p. 72 pl. xxx figs. 5-8, armiger pp. 71-72 pl. xxx figs. 13-15, Cooke.

†Periarchus p. 41 with key to spp. from eastern U.S., P. lyelli pp. 41-42 pl. xiii fig. 1 pl. xiv figs. 1-3, lyelli floridanus p. 42 pl. xiii fig. 4, lyelli pileus-sinensis p. 42 pl. xiii figs. 2, 3, protuberans pp. 42-43 pl. xii figs. 7-9, quinquefarius pp. 43-44 pl. xiv figs. 6-8, quinquefarius kewi p. 44 pl. xiv figs. 4, 5, Cooke.

Peronella japonica p. 117 pl. lix fig. 5, minuta p. 117 pl. lix fig. 4, Utinomi.

Phyllacanthus parvispinus, Sinclair p. 4 fig.

†Phyllacanthus pp. 10-11 with P. mortoni pp. 11-12 pl. i figs. 15-17, Cooke.

†Phymosoma ornatissimum, possible identity of Rumphius', 1705, pl. lix fig. 6, Engel p. 222 photo 23 fig. 5.

†Phymotaxis p. 24, with P. mansfieldi p. 24 pl. vi figs. 4-8, Cooke.

Plagiobrissus grandis, Cooke pp. 86–87 pl. xxxix figs. 1, 2 pl. xl fig. 6, Tommasi p. 7 pl. i fig. 4.

†Plagiobrissus p. 86, with P. curvus pp. 87-88 pl. xxxix figs. 3-6, P. ? dixie p. 87 pl. xl figs. 1-5, P. ? holmesii p. 87, Cooke.

†Porpitella p. 33, with P. micra p. 33 pl. vii figs. 5-7, Cooke.

Prionocidaris baculosa var. annulifera, Utinomi p. 113 pl. lvii fig. 1.

†Protoscutella p. 38, with P. conradi pp. 39-40 pl. xv figs. 4, 5, mississippiensis pp. 38-39 pl. xv figs. 12, 13, pentagonium p. 39 pl. xv figs. 6-8, plana p. 40 pl. xv figs. 1-3, tuomeyi p. 39 pl. xv figs. 9-11, Cooke.

†Psammechinus p. 16 with P. philanthropus pp. 16-17 pl. iii figs. 1-2, santee p. 17 pl. iii figs. 3-4, Cooke; P. cf. salvensis, Mitrović- Petrović p. 48 pl. i fig. 3.

Pseudocentrotus depressus, Utinomi p. 115, pl. lviii fig.

†Pseudodiadema jacardi pp. 47-48 pl. i fig. 1, raulini pp. 46-47 pl. i fig. 2, Mitrović-Petrović.

†Pygaulus desmoulinsii p. 53 pl. iii fig. 2, ovatus p. 54 pl. iii fig. 3, Mitrović-Petrović.

†Pygomalus analis pp. 340-341 fig. 1A, ovalis p. 340 pl. i figs. 1, 2, Jesionek-Szymanska.

†Pygorhytis ringens, Jesionek-Szymanska p. 341 fig. 1B.

†Pyrina cylindrica p. 51 pl. ii fig. 3, loevis pp. 52-53 pl. iii fig. 1, pygea pp. 51-52 pl. ii fig. 4, Mitrović-Petrović.

†Rhopostoma gen. nov. family Echinoneidae, for type Ananchytes cruciferus Morton 1830, p. 26, R. cruciferum p. 26 pl. vii figs. 1-4, Cooke.

Rutula orbiculus, Cherbonnier (3) p. 49 pl. vii fig. K; R. orbiculus, identity of Rumphius' pl. xiv fig. 1, Engel p. 217 photo 24 fig. 5.

†Salenia p. 13 with S. tumidula p. 13 pl. ii figs. 1-7, Cooke.

Salmacis bicolor, Halstead fig. 25 (pedicellariae); S. sphaeroides, Gillett & McNeill pl. xcv fig. 2.

†Santeelampas gen. nov. family Cassidulidae, for type Catopygus oviformis Conrad 1850, p. 61, S. oviformis pp. 61-62 pl. xxvi figs. 1-8, Cooke.

Scaphechinus brevis p. 118 pl. lix fig. 7, mirabilis p. 118 pl. lix fig. 8, tenuis pp. 117-118 pl. lix fig. 6, Utinomi.

Schizaster canaliferus, Tortonese (1) p. 292; S. edwardsi, Cherbonnier (3) pp. 50-51 pl. viii figs. F-L; S. lacunosus, identity of Rumphius', 1705, pl. xiv fig. 2, Engel pp. 215-216 photo 24 fig. 12; S. lacunosus, Utinomi p. 119 pl. lx fig. 4.

†Schizaster p. 72, with S. (Brachybrissus) p. 73, S. (Brachybrissus) ocalanus p. 73 pl. xxix figs. 13-17, Cooke.

Spatangus purpureus, Cherbonnier (3) p. 50 pl. viii figs. A-E, Tortonese (1) p. 292, Nichols (2) pp. 353-354 363-368, 374-375 figs. la-c, 4a-c, 7a-c, 12, 14, 15 (part), 20, 31, raschi pp. 355, 373-374, 376-377 figs. 2a-c, 5a-c, 8a-c.

†Spatangus p. 80, with S. glenni sp. nov. Miocene S. Carolina, U.S. p. 80 pl. xxxv figs. 1-5, Cooke.

Sphaerechinus granularis, Sebastio p. 100 figs. 5, 6. Stomopneustes variolaris, Utinomi p. 113 pl. lvii fig. 3. Strongylocentrotus p. 25, with S. dröbachiensis p. 25 pl. vi fig. 3, Cooke, also Madsen (2) p. 169 fig. 5 (holotype), Scheffer pp. 373-374 fig. 5; S. pallidus, Tambs-Lyche pp. 10-11.

Stylocidaris affinis, Tortonese (1) p. 291; S. lineata, Cherbonnier (1) p. 367.

Temnopleurus reevesi p. 114 pl. lvii fig. 11, toreumaticus p. 114 pl. lvii fig. 10, Utinomi.

Temnotrema sculptum, Utinomi p. 114 pl. lvii fig. 9. †Trematopygus crucifer (Morton) made the type of a new genus Rhopostoma by Cooke p. 26.

Tripneustes p. 15 with T. ventricosus pp. 15–16 pl. ii figs. 14–16, Cooke; T. gratilla, Alcala pl. i; T. gratilla, identity of 'Echinus esculentus' of Rumphius 1705 p. 31, I, Engel pp. 210–211; T. gratilla, Gillett & McNeill pl. cliii, Tahara, Okada & Kobayashi pp. 168–169 fig. 3 pl. vii figs. 5, 6, Utinomi p. 115 pl. lviii fig. 2.

†Toxaster amplus, Kotetishvili pp. 20-21 pl. i figs. 2a-c.

Toxopneustes chlorocanthus, Alcala pl. iii; T. elegans, Halstead fig. 23; T. pileolus, Tahara, Okada & Kobayashi p. 168 pl. viii figs. 3, 4, Utinomi p. 115 pl. lviii fig. 1.

 $\dagger Tylocidaris macneili pp. 12–13 pl. i figs. 3–5, <math display="inline">T.$? salina p. 12 pl. i figs. 12–14, both spp. nov. Paleocene Alabama, U.S., T. walcotti p. 12 pl. i figs. 7–9, Cooke.

†Unifascia gen. nov. family Unifasciidae, for type Macropneustes carolinensis Clark 1915, pp. 79-80, U. carolinensis p. 80 pl. xxxiv figs. 7-13, Cooke.

Unifasciidae fam. nov., presumably order Spatangoida, for genus *Unifascia* nov., **Cooke** p. 79.

†Weisbordella p. 53, with W. cubae pp. 53-54 pl. xx figs. 1-4, johnsoni pp. 54-55 pl. xx figs. 5-7, Cooke.

 $\dagger Wythella$ pp. 52–53, with W. eldridgei p. 53 pl. xxi figs. 8–10, Cooke.

†SOMASTEROIDEA

[Vacant]

ASTEROIDEA

Acanthaster planci, identity of 'Stella marina quindecim radiorum' of Rumphius, 1705, Engel pp. 218-219; A. planci, Gillett & McNeill pl. lxxxix (coloured), Utinomi p. 110 pl. ly fig. 7.

Allostichaster insignis p. 142 figs. 39, 40, polyplax p. 142 fig. 38, Fell.

Anareaster gen. nov., family Korethrasteridae, for type A. ganymede sp. nov. MacRobertson Land, Antarctica, Fell & Clark, H. E. pp. 185-186 figs. 1-3.

Anthenoides granulosus, Fell p. 135.

Asterias amurensis, Scheffer p. 372 fig. 3, Utinomi p. 110 pl. lv fig. 9.

Asterina aucklandensis p. 138 fig. 23A, novaezelandiae p. 138, regularis p. 138 fig. 23, Fell; A. batheri p. 109 pl. lv fig. 4, coronata japonica p. 109 pl. lv fig. 3, pectinifera pp. 109–110 pl. lv fig. 5, Utinomi.

Asterodiscus truncatus, Fell p. 135 fig. 17.

Asterodon dilatatus p. 134 fig. 10, miliaris p. 134 fig. 8, robustus p. 134 figs. 9, 14, Fell.

Astropecten duplicatus pp. 108-109 fig. 1, richardi pp. 110-111 fig. 2, Cherbonnier (1); A. dubiosus, polyacanthus p. 131, primipenius p. 131 fig. 6, Fell; A. aranciacus, Madsen (2) pp. 165, 167, notes on the holotype; A. armatus brasiliensis pp. 12-13 pl. ii fig. 3,

armatus riensis pp. 13–14 pl. ii fig. 4, marginatus p. 14 pl. ii fig. 5, Tommasi; A. irregularis pentacanthus, Tortonese (1) p. 290; A. latespinosus p. 107 pl. liv fig. 3, polyacanthus p. 107 pl. liv fig. 2, scoparius p. 107 pl. liv fig. 1, Utinomi.

Astrostole scabra, Fell p. 140 fig. 41.

Benthopecten pentacanthus, Fell p. 134.

Brisingines delli, Fell p. 140 fig. 33.

Calvasterias laevigata p. 142, suteri p. 142 fig. 42, Fell'

Certonardoa semiregularis, Utinomi p. 109 pl. lv fig. 1.

Cheiraster richardsoni, Fell p. 134 fig. 30.

Chondraster a subgenus of Porania with C. hermanni sp. nov. East Greenland, Madsen (1) pp. 153-160 figs. 1-4, C. elattosis probably not specifically distinct from grandis pp. 156, 157.

Coscinasterias calamaria, Fell p. 142 fig. 37; C. acutispina, Utinomi p. 110 pl. lv fig. 8.

Cosmasterias dyscrita, Fell p. 142 figs. 31, 35.

Crossaster japonicus, Fell p. 139 fig. 27.

Culcita novaeguineae, Gillett & McNeill pl. lxxxix (coloured), Utinomi p. 108 pl. liv fig. 9.

Dipsacaster magnificus, Fell p. 132 fig. 3.

Echinaster echinophorus, Cherbonnier (1) pp. 171–172 fig. 3C, D; E. farquhari, Fell p. 139 fig. 24; E. antonioensis p. 22 pl. iv fig. 2, brasiliensis pp. 22–23 pl. iv fig. 3, sentus pp. 23–24 pl. iv fig. 4, spinulosus pp. 21–22 pl. iv fig. 1, Tommasi; E. sepositus, Tortonese (1) p. 291.

Eurygonias hylacanthus, Fell pp. 134-135 figs. 11, 15. Fromia sp., Gillett & McNeill pl. lxxxviii.

Goniaster americanus, Cherbonnier (1) pp. 107-108.

Goniodiscaster scaber with synonym Asterias articulata Linnaeus 1753, **Madsen** (2) pp. 165-167 fig. 2 (type of articulata).

Hacelia attenuata, Tortonese (1) p. 291.

Henricia compacta etc. p. 139 fig. 25, ralphae p. 139 fig. 26, Fell; H. ohshimai, Utinomi p. 110 pl. lv fig. 6.

Hippasteria trojana, Fell, pp. 136, 138 fig. 21.

†Hudsonaster wardi sp. nov. Devonian Pennsylvania, U.S., Cramer, pp. 471-473 fig. 1.

Leiaster leachi, Utinomi p. 109 pl. lv fig. 2.

Linckia laevigata, identity of 'common species' of Rumphius, 1705 p. 39 and pl. xv fig. E, Engel p. 218 photo 25 fig. 2; L. laevigata, Gillett & McNeill pl. xc (coloured); L. guildingii, Tommasi pp. 17-18 pl. iii fig. 3; L. guildingi, Utinomi p. 108 pl. liv fig. 6.

Luidia barbadensis pp. 170-171 fig. 3 B, E, fig. 4, clathrata p. 170, senegalensis pp. 169-170, Cherbonnier (1); L. neozelanica p. 132 fig. 7, varia p. 132, Fell; L. clathrata p. 9 pl. ii fig. 1, senegalensis pp. 9-11 pl. ii fig. 2, Tommasi; L. maculata p. 108 pl. liv fig. 7, quinaria p. 108 pl. liv fig. 8, Utinomi.

Mediaster sladeni, Fell p. 136 fig. 19.

Nardoa frianti, Utinomi p. 108 pl. liv fig. 5.

Nectria pedicelligera, Fell p. 135 fig. 18.

Odontaster benhami, Fell p. 135.

Ophidiaster confertus, Gillett & McNeill pl. cliii; O. kermadecensis, Fell p. 135 fig. 16; O. cribrarius, Utinomi p. 107 pl. liv fig. 4.

Oreaster clavatus with synonyms Asterias dorsata Linnaeus 1753 and A. nodosa Linnaeus 1758 pp. 163–164, O. reticulatus with synonym Asterias gigas Linnaeus 1753 p. 163 fig. 1 (type of gigas), O. turritus Müller & Troschel valid for the species commonly known as nodosus Linnaeus pp. 164–165, Madsen (2); O. reticulatus, Tommasi pp. 16–17 pl. iii fig. 2.

Patiria stellifer, Tommasi pp. 19-20 pl. iii fig. 4.

Pentaceraster alveolatus or multispinus, identity of Rumphius', 1705 pl. xv fig. D, Engel p. 219 photo 25 fig. 1.

Pentagonaster pulchellus, Fell p. 136 fig. 20.

Peribolaster lictor, Fell p. 139 fig. 29.

Persephonaster neozelanicus, Fell pp. 131-132 fig. 5.

Plutonaster knoxi, Fell p. 132.

Porania (Chondraster) hermanni sp. nov. Madsen (1), see under Chondraster; P. pulvillus, Madsen (2) pp. 169-170 fig. 6 (holotype).

Poraniidae a synonym of Asteropidae, Madsen (1) p. 155.

Protoceaster nodosus, identity of 'Stella Marina Quarta' or 'Artocreas Marinum' of Rumphius, 1705 pl. xv fig. A, Engel p. 219 photo 25 fig. 3.

Pseudarchaster abernethyi p. 136 fig. 4, garricki p. 136, Fell.

Pseudoporania stormi, Madsen (1) pp. 157, 158, referred to Porania (Chondraster).

Psilaster acuminatus, Fell p. 131 fig. 13.

Pteraster bathami, Fell p. 139 fig. 28.

Pycnopodia helianthoides, Scheffer p. 373 fig. 4.

Sclerasterias mollis, Fell p. 140 fig. 32; S. richardi, Tortonese (1) p. 291.

Solaster endeca, identity of Rumphius', 1705 pl. xv fig. F, Engel p. 219 photo 25 fig. 4.

Sphaeriaster berthae, björlykkei, Madsen (1) p. 158, referred to Porania (Chondraster).

Stegnaster inflatus, Fell, p. 138 fig. 22.

Stichaster australis, Fell p. 142 fig. 34.

Tethyaster vestitus, Cherbonnier (1) p. 168 fig. 3A, Tommasi pp. 15-16 pl. iii fig. 1.

Zoroaster spinulosus, Fell p. 140 fig. 36.

OPHIUROIDEA

Amphiodia occidentalis, Light and others fig. 123b.

Amphiura griegi, Tambs-Lyche p. 11; A. apicula
pp. 293-294, chiajei, filiformis p. 293, Tortonese (1).

Astrophyton muricatum, Cherbonnier (1) p. 261.

Gorgonocephalus caryi, Utinomi p. 111 pl. lvi fig. 1. Hemipholis elongata, Parker pl. iii fig. 15a-c.

Macrophiothrix longipeda, Gillett & McNeill, pl. xciii fig. 7, Utinomi pp. 111-112 pl. lvi fig. 5.

Ophiactis savignyi, Utinomi p. 111 pl. lvi fig. 3.

Ophiarachna incrassata, Gillett & McNeill pl. xcii figs. 1, 2.

Ophiarachnella infernalis, Gillett & McNeill pl. xcii, fig. 3; O. gorgonia, Utinomi p. 112 pl. lvi fig. 10.

Ophiarthrum pictum, Gillett & McNeill pl. xeiia fig. 6.

Ophiocoma erinaceus, insularia, Gillett & McNeill pl. xcii figs. 4, 5; O. brevipes, Utinomi p. 112 pl. lvi fig. 8.

Ophioderma brevispina, Cherbonnier (1) pp. 261-262 fig. 5A, B fig. 6A, B.

Ophiolepis elegans, Cherbonnier (1) p. 263, Parker pl. iii fig. 14a-c.

Ophiomastix annulosa p. 112 pl. lvi fig. 6, mixta p. 112 pl. lvi fig. 7, Utinomi.

Ophiomyxa brevispina, possible identity of 'Stella marina quinta etc. scolopendroides' of Rumphius, 1705, p. 40 and pl. xv figs. B, C, Engel p. 220 photo 26 figs. 3, 2; O. pentagona, Tortoness (1) p. 292.

Ophioncus granulosus, Light & others fig. 123 f.

Ophiopholis aculeata, Light & others fig. 123a, c.

Ophioplocus esmarki, Light & others fig. 123d; O. japonicus, Utinomi p. 112 pl. lvi fig. 9.

Ophiopsila annulosa, Tortonese (1) p. 294.

Ophiothrix spiculata, Light & others fig. 123g; O. fragilis, Tortoness (2) pp. 1-9.

Ophiotrichoides nereidina, Utinomi p. 111 pl. lvi fig. 4.

Ophiura fallax sp. nov. French Guiana, Cherbonnier
(1) pp. 263-265 fig. 7A-D.

†Ophiura graysonensis, Clark, D. L. pp. 1126-1127.

†Ophiuraster burrisi, Hattin pp. 1125-1126 fig. 1.

Trichaster elegans, Utinomi p. 111 pl. lvi fig. 2.

Indeterminable Euryalid, 'Caput medusae' of Rumphius, 1705, p. 41 pl. xvi figs. Å, B, Engel pp. 220–221 photo 26 figs. 1, 2.

CRINOIDEA

†Acanthocrinus p. 125 with A. aff. jaekeli pp. 125-127 pl. ii figs 1-3, Le Maître.

Antedon mediterranea, Tortonese (1) p. 290,

†Codiacrinidae, Le Maître pp. 141-142 text-fig. 3 pl. ii fig. 4.

†Collarocrinus subgen, nov. of Torynocrinus, for type C. phialaeformis sp. nov., Szörényi p. 250.

Comantheria grandicalyx, Utinomi p. 105 (in Japanese) pl. liii fig. 2.

Comanthus parvicirra, Gillett & McNeill pl. xci (coloured); C. (Cenolia) japonica pp. 105-106 pl. liii fig. 3, C. (Comanthus) parvicirra p. 106 pl. liii fig. 4, Utinomi,

†Ctenocrinus, Le Maître p. 131 pl. ii fig. 5.

†Cyclocyclicus brevitodentatus Carboniferous pp. 47, 65-66 pl. i fig. 14, chaneensis pp. 53, 73 pl. ii fig. 6, disparies pp. 56, 76 pl. ii fig. 10, both Permian, kuangtungensis Carboniferous pp. 52-53, 72-73 pl. ii fig. 5, kueichounensis Triassic pp. 47, 66 pl. i fig. 15, mui Permian pp. 54, 56, 75 pl. ii fig. 9, perpusilus Carboniferous pp. 47, 65 pl. i fig. 13, schizinicus Triassic pp. 48, 66-67 pl. i fig. 18, suni pp. 51-52, 71-72 pl. ii fig. 4, tiens pp. 49-50, 69-70 pl. ii fig. 2, both Carboniferous, all spp. nov. from S. China, Dubatolowa & Shao, also C. circunvalatus pp. 48-49, 67-69 pl. ii fig. 1, mirandus pp. 46-47, 64-65 pl. i fig. 12, orenarius var. ordinata pp. 54, 74-75 pl. ii fig. 8, pulcher pp. 50-51, 70 pl. ii fig. 3, C. aff. dignus pp. 53-54, 73-74 pl. ii fig. 7, all Carboniferous S. China, Yeltyschewa in Dubatolowa & Shao (given as 'in litt.', not previously recorded but ? nov., Rec.); C. heui pp. 48, 66 pl. i fig. 17, liliformis pp. 48, 67 pl. i fig. 19, 16.

†Cyrtocrinus remeši Sp. nov. Cretaceous Hungary (†Rec.), for C. marginatus (pars) Remeš 1902 p. 11 pl. xviii figs. 21, 22, 24, Szörényi pp. 239–240.

Decametra tigrina, Utinomi p. 106 pl. liii fig. 5.

† Edriocrinus pp. 142-143 with E. cf. pocilliformis pp. 143-145 pl. ii figs. 7-9, Le Mattre.

†Gastrocrinus p. 133 with G. patulus pp. 133-134 pl. ii fig. 11, Le Maître.

†Griphocrinus p. 127 with G. nodulosus p. 128, Le Maitre.

†Haplocrinites p. 132 with H. aff. clio pp. 132–133 pl. iii fig. 7, Le Maître.

†Labiocrinus subgen. nov. of Torynocrinus, for type L. labiatus sp. nov., Szörényi pp. 253-254.

†Lecanocrinidae?, Le Maître pp. 145-146 pl. i figs. 11-13.

Leptometra phalangium, Tortonese (1) p. 290.

†Pentagonocyclicus admotus Permian S. China pp. 43-44, 60 pl. i fig. 5, expressus pp. 43, 59-60 pl. i fig. 4, falsus pp. 45, 63 pl. i fig. 10, hsianghsiangensis pp. 46, 64 pl. i fig. 16, kuangsinensis pp. 43, 59, pl. i fig. 3, pulverlus pp. 42-43, 58 pl. i fig. 2, schansinicus pp. 45-46, 63-64 pl. i fig. 11, stellatus pp. 44, 60-61 pl. i fig. 6, triformis pp. 44-45, 61-62 pl. i fig. 8, all Carboniferous S. China, spp. nov., Dubatolowa & Shao, P. circumvalatus var. minor pp. 45, 62 pl. i fig. 9, dvinae-boreae pp. 44, 61 pl. i fig. 7, both Yeltyschewa in Dubatolowa & Shao (given as 'in litt.', not previously recorded but ? nov., REC.)

†Pentececrinus gen. nov. suborder Cyathocrinoidea, pp. 464-467, for type P. parvus sp. nov. Devonian Missouri, U.S., pp. 467-469 text-figs. 1, 3a-g, Koenig & Niewoelner.

†Phyllocrinus pp. 151-152 with P. furcillatus sp. nov. Jurassic New Zealand, Speden pp. 152-154 pl. xx.

†Semiometra italica, Balavoine p. 525.

† Thylacocrinus pp. 128–129 with T. aff. vannioti pp. 129–131 pl. ii fig. 6 pl. iii fig. 6, **Le Maître**.

†Tiaracrinus pp. 134-135 with T. quadrifrons pp. 135-139 pl. i figs. 1-5, soyei pp. 139-140 pl. i figs. 8-10, Le Maître.

†Torynocrinus pp. 232–236 with Torynocrinus pp. 240–241, Collarocrinus p. 250, Labiocrinus pp. 253–254, subgenera nov., T. (Torynocrinus) bellus pp. 246–247 pl. ii figs. 29–23, compactus pp. 247–249 pl. ii figs. 1–8, floriformis pp. 245–246 pl. i. figs. 29–32, hungaricus pp. 243–245 pl. i figs. 1–24, sulcatus pp. 249–250 pl. ii figs. 9–19, spp. nov. Cretaceous Hungary, T. (Torynocrinus) pranulatus pp. 241–243 pl. i figs. 25–28, T. (Collarocrinus) phialaeformis pp. 251–252 pl. iii figs. 1–4, pulcher pp. 252–253 pl. iii figs. 5–8, both spp. nov. Cretaceous Hungary, T. (Labiocrinus) labiatus pp. 254–256 pl. iii figs. 9–15, minor pp. 256–257 pl. iv figs. 1–8, both spp. nov. Cretaceous Hungary, T. maryinatus pp. 238–239, thersites pp. 236–237, thersites var. difformis p. 238, thersites var. palmata p. 237, Szőrényi.

†Trigonotrigonalis asymmetricus Yeltyschewa in Dubatolowa & Shao pp. 42, 57-58 pl. i fig. 1 (given as 'in litt.', not previously recorded but ? nov., Rec.)

Tropiometra carinata, Cherbonnier (1) p. 107; T. afra macrodiscus, Utinomi p. 105 (in Japanese) pl. liii fig. 1.

†Crinoidea, Zakowa p. 126 pl. x figs. 4a-e.

†MACHAERIDEA

[Vacant]

†EOCRINOIDEA

[Vacant]

†PARACRINOIDEA

[Vacant]

†CYSTOIDEA

Eucystis pp. 120-121 with E. sp. pp. 121-122 text-figs. 1, 2 pl. iii figs. 4, 5, Le Maître.

Heliocrinites ?, Le Maitre pp. 118-120 pl. iii figs. 1-3.

†CARPOIDEA

†Cincta or Trochocystids, Cabibel, Termier & Termier pp. 287, 288.

Decacystis, Cabibel, Termier & Termier pp. 290, 291.

†Eikosacystis gen. nov. aff. Carpoid Cincta, p. 286 for type E. couloumanensis sp. nov. text-fig. 10 pp. 286, 287 text-fig. 5 pl. i fig. 5, E. ? ferralsensis sp. nov. p. 287 pl. i figs. 6, 7, both Cambrian S. France, Cabibel, Termier & Termier.

Gyrocystis pp. 289, 290 with G. platessa p. 290 textfig. 7, Cabibel, Termier & Termier.

Trochocystites p. 288 with T. bohemicus p. 289 text-fig. 6 pl. ii figs. 1–6 (T. barrandei Munier-Chalmas & Bergeron 1889 a synonym), Cabibel, Termier & Termier.

†EDRIOASTEROIDEA

[Vacant]

†BLASTOIDEA

Pentremitidea p. 123 with P. cf. pailletti pp. 123-124 pl. ii fig. 10, Le Maître.

INCERTAE SEDIS

†Cambraster p. 284 with C. cannati pp. 284, 285 text-figs. 3, 4 pl. i figs. 2-4, Cabibel, Termier & Termier.

†Sucocystis gen. nov. for type S. theronensi sp. nov. Cambrian S. France, Cabibel, Termier & Termier p. 291 text-fig. 9, pl. ii figs. 7-9.

†Stromatocystites pp. 283, 284 with S. sp. p. 284 pl. i fig. 1, S. pentangularis text-fig. 2, Cabibel, Termier & Termier.

THE ZOOLOGICAL RECORD

THE object of The Zoological Record is to give, by means of an annual volume, complete lists of the works and publications relating to zoology in all its branches that have appeared during the year preceding the issue of the volume, together with a subject and systematic index.

The complete volume can be obtained from the Society at the price of £10 0s. Od. per copy.

The following volumes are available:-

In Reprint Edition

Vols, 1-30 (1864-1893) obtainable only from Messrs. Butterworths, 4-5, Bell Yard, London, W.C.2.

Complete set of the first 25 volumes, bound in cloth, price £221 0s. 0d.

Complete set of 30 volumes, in paper covers, price £256 10s. 0d.

Separate volumes, in paper covers, vols. 1-20 £8 0s. 0d. each; Vols. 21-25 £8 18s. 6d. each; Vols. 26-30 £10 10s. 0d. each.

In Original Edition

Vols. 65-71 (1928-1934) at £5 each.

Vols. 85-91 (1948-1954) at £6 each.

Vols. 92-94 (1955-1957) at £8 each.

For particulars of the availability and prices of separate Sections, application should be made to the Scientific Director, The Zoological Society of London, Regent's Park, London, N.W.1.

It should be noted that the separate Section Insects can only be obtained from the Commonwealth Institute of Entomology, 56 Queen's Gate, London, S.W.7.

NOMENCLATOR ZOOLOGICUS.—Lists the names of all the genera and subgenera in zoology from the 10th Edition of Linnaeus 1758 to the end of 1945, with a bibliographical reference to the original description of each. Price 17½ guineas, postage extra. Price of separate volumes: Vols. 1-4 (1758–1935): Vol. 5 (1936–1945) 3½ guineas each, postage extra.

SEPARATE SECTIONS OF THE ZOOLOGICAL RECORD

In addition to the complete bound volume, the separate Sections may be obtained singly, bound in printed paper covers.

The prices of	the Sections from	m Vo	lume 9	6 are :-				ī		
1.	Comprehensive	Zoolo	ogy			•••			s. 4	d. 0
2.	Protozoa	•••			•••		***		13	0
3.	Porifera	•••				•••			3	0
4.	Coelenterata	•••	***	•••	***	•••	***	•••	5	0
5.	Echinodermata		***	•••	•••	•••	•••	•••	4	0
6.	Vermes		•••	•••	***			•••	12	0
7.	Brachiopoda	•••		***	***	•••	•••		4	0
8.	Bryozoa	•••	***	***			***	•••	3	0
9.	Mollusca	***	•••	***	•••	•••		•••	16	0
10.	Crustacea	•••	•••	***	***	.***	***	•••	8	0
11.	Trilobita	•••		•••	•••		•••	•••	8	0
12.	Arachnida	•••	•••	•••	•••			• • •	14	0
13.	*Insecta	•••		•••		•••	•••	•••	60	0
14.	Protochordata	•••	•••	•••	•••	•••	•••		3	0
15.	Pisces	•••	•••	•••	•••	•••	•••	•••	11	0
16.	Amphibia	•••	•••	•••	•••	•••	***	•••	. 12	0
17.	Reptilia	***	•••	•••	•••	***		•••	12	0
18.	Aves	•••	•••	•••	•••	•••		•••	12	0
19.	Mammalia	•••	•••	***	•••	•••	•••	•••	12	0
20.	List of New Genera and Subgenera					***	•••	•••	4	0

 $[\]boldsymbol{*}$ Obtainable only from the Commonwealth Institute of Entomology, 56 Queen's Gate, London, S.W.7.

